


Inclusive b production at LHC with CMS

Valery Andreev


UCLA

June 8, 2006


CERN, 2nd HERA-LHC workshop



Large Hadron Collider (LHC)



The Large Hadron Collider (LHC)




- Design luminosity $L = 10^{34} \text{ cm}^{-2} \text{ s}^{-1}$
 $\sim 100 \text{ fb}^{-1} / \text{year}$
 Pile up ~ 20 collisions/crossing
 40 MHz pp bunch-crossing rate

- Start-up luminosity $L \approx 10^{33} \text{ cm}^{-2} \text{ s}^{-1}$
 $\Rightarrow \sim 10 \text{ fb}^{-1} / \text{year}$

- expected completion : mid 2007

	Beams	Energy	Luminosity
LEP	$e^+ e^-$	200 GeV	$10^{32} \text{ cm}^{-2} \text{ s}^{-1}$
LHC	$p p$	14 TeV	$10^{34} \text{ cm}^{-2} \text{ s}^{-1}$
	$Pb Pb$	1312 TeV	$10^{27} \text{ cm}^{-2} \text{ s}^{-1}$



The CMS detector

The Compact Muon Solenoid (CMS)

Onion structure


- Tracker
- Calorimeters
- Muon system

Precise


$e, \mu, \gamma, \text{jets}, E_T$

Efficient

b tagging, τ detection



B production



- *b production* at hadron colliders
 - Huge cross section
 - Challenge for perturbative QCD
 - New physics searches: b jets as a signal feature

Analysis

- **B production total cross section**
- **Differential cross sections $d\sigma/dp_t$, $d\sigma/d\eta$**
 - **Selection (b-tag)**
 - **semileptonic b-decays into muons**
 - **Background (b purity)**
 - **Trigger efficiency**
 - **Luminosity**

Event Selection

B tagged jet + muon

- Trigger
 - Level 1: single muon, $P_t > 14 \text{ GeV}/c$
 - High Level (HLT) trigger, cross-channel
 - Muon, $P_t > 19 \text{ GeV}/c$, $|\eta| < 2.4$
 - B tagged Jet, $E_t > 50 \text{ GeV}$
- B tag
 - Inclusive secondary vertex reconstruction in jets (C. Weiser, CMS Note 2006/014, 2006)

B phase space



- *B* hadron
 - $P_t > 50 \text{ GeV}/c$
 - $|\eta| < 2.4$

Event selection



~4 M QCD events processed

\hat{p}_T , GeV/c	σ^{QCD} , μb	$N_{\text{generated}}^{\text{QCD}}$, events	$b\bar{b}$ purity, %	$c\bar{c}$ fraction, %	uds fraction, %	$N_{\text{expected}}^{b\bar{b}}$, events
50 – 80	20.9	198993	66	32	2	1.4 M
80 – 120	3.0	294986	66	32	2	6.1 M
120 – 170	0.5	291982	72	26	2	5.1 M
170 – 230	0.1	355978	71	26	3	2.4 M
230 – 300	2.4×10^{-2}	389978	73	24	3	0.9 M
300 – 380	6.4×10^{-3}	283983	70	25	5	0.3 M
380 – 470	1.9×10^{-3}	191989	68	27	5	88 k
470 – 600	6.9×10^{-4}	190987	64	29	7	34 k
600 – 800	2.0×10^{-4}	94996	60	31	9	10 k
800 – 1000	3.6×10^{-5}	89999	60	30	10	2.0 k
1000 – 1400	1.1×10^{-5}	89998	55	31	14	0.5 k



Level 1 trigger efficiency



High Level (HLT) trigger efficiency




B tagging efficiency







B jet resolution

$P_t > 170 \text{ GeV}/c$





B tagging with muon



Muon is inside jets



B tagging with muon



B jet-muon association efficiency




~ 75 %

Fit results

QCD events MC: $120 < P_t < 170 \text{ GeV}/c$

Muon P_t w.r.t. the closest B jet



Nb = 2503 (66%)
 Nc = 965 (26%)
 Nuds = 299 (8%)

- b -

- c -

3767 events

- uds -

Fit:


Nb = 2750 ± 346
 Nc = 702 ± 513
 Nuds = 329 ± 235

 3781 events

Fit results

QCD events MC: $230 < P_t < 300 \text{ GeV}/c$

Muon P_t w.r.t. the closest B jet



	Nb	=5250 (56 %)
	Nc	= 2388 (26%)
\underline{b} - - -	Nudsg=	1740(18%
\underline{c} . . .		-----
udsg		9378 events
.....		

Fit:

Nb	=5222 ± 501
Nc	= 2050 ± 728
Nudsg	=1778 ±341

	9050 events

Systematics ($\geq 10 \text{ fb}^{-1}$)

Source	uncertainty, %
jet energy scale	12
event selection	6
B tagging	5
luminosity	5
trigger	3
muon Br	2.6
misalignment	2
muon efficiency	1
$t\bar{t}$ background	0.7
fragmentation	9
total	18

Conclusions

- **~16 M b events to be selected at 10 fb^{-1} by CMS (one year of low lumi LHC)**
- **Up to 1.5 TeV P_t reach is expected**
- **New test of QCD is coming**