

Higgs production at the LHC – Theory Status –

Robert Harlander

CERN, July 8, 2003

Introduction

Gluon fusion


Weak Boson Fusion

Higgs Strahlung


$t\bar{t}H$ & $b\bar{b}H$

Conclusions and Goals


Production Modes




Gluon fusion



Weak Boson Fusion





Higgs Strahlung




$t\bar{t}H$

Cross sections at the LHC



Combination: Couplings




[Duehrssen, Jakobs]

→ talk by [D. Zeppenfeld](#)

Thu, 9.00am, Higgs session, Council Chamber


Diffractive Higgs Production




- protons detected in roman pot detectors
- first investigations ('90):
[Bialas, Landshoff, Szeremeta, Janik, Nachtmann, Schäfer, Schöpf, ...]
- recent activity: → talk by [A. de Roeck](#)

Wed 13.45, Higgs Session, Council Chamber


Production Modes




Gluon fusion



Weak Boson Fusion




Higgs Strahlung




$t\bar{t}H$

Higgs discovery potential



Total cross section for $gg \rightarrow H + X$



- see also [Ravindran, Smith, v.Neerven ('03)]
- soft gluon resummation: +6% [Catani, de Florian, Grazzini, Nason]

Distributions

- Higgs Rapidity and Transverse Momentum at NLO:

$$\frac{d^2\sigma}{dp_T dy}, \quad \frac{d\sigma}{dp_T}, \quad \frac{d\sigma}{dy}$$

[de Florian, Grazzini, Kunszt ('99)]

[Glosser, Schmidt ('02)]

[Ravindran, Smith, v.Neerven ('02)]

[Bozzi, Catani, de Florian, Grazzini ('03)]

[Anastasiou, Dixon, Melnikov ('03)]

→ talk by [M. Grazzini](#)

Wed, 2.10pm, Higgs session, Council Chamber


$gg \rightarrow H$: Open questions

- effective Lagrangian valid at NNLO?
- NNLO parton densities?
- Tools? NNLO Monte Carlo?
- K-factors in experimental analyses?

→ talk by [B. Mellado](#)


Wed, 16.45, Higgs Session, Council Chamber

- K-factors for distributions:




[de Florian,
Grazzini,
Kunszt ('99)]


Production Modes




Gluon fusion



Weak Boson Fusion




Higgs Strahlung




$t\bar{t}H$

Weak Boson Fusion




- [Rainwater, Zeppenfeld, Plehn, Kauer, Hagiwara, Eboli,...]
- two forward jets
- helpful for $M_H < 140 \text{ GeV}$
- couplings, CP, ...




[Mangano, Moretti, Piccinini, Pittau, Polosa ('03)]

Higgs discovery potential



Weak Boson Fusion at NLO



~ Deep inelastic scattering!
[Han, Valencia, Willenbrock ('92)]

$\approx +10\%$


jet distributions [Figy, Oleari, Zeppenfeld ('03)]

→ Tools?


→ talk by [C. Oleari](#)

Wed, 14.35, Higgs Session, Council Chamber


Production Modes




Gluon fusion



Weak Boson Fusion






Higgs Strahlung




$t\bar{t}H$

Higgs Strahlung




- most important for Tevatron
- **relevance for LHC?**




?

Higgs Strahlung at NNLO


[R.H., Djouadi (in prep)]




Production Modes




Gluon fusion



Weak Boson Fusion




Higgs Strahlung



$t\bar{t}H$

Associated $t\bar{t}H$




- small rate but clear signature
- top Yukawa coupling
- **NLO** prediction very stable
- Tools? Also useful for $b\bar{b}H$?

→ talk by [M. Spira](#)

Wed, 3.15pm, Higgs Session, Council Chamber

$t\bar{t}H$: LHC



[Beenakker, Dittmaier, Krämer, Plümper, Spira, Zerwas ('01)]

[Dawson, Reina, Wackerth, Orr, Jackson ('03)]


Supersymmetry

- **charged Higgs:** → talk by **T. Plehn**
Wed, 3.00pm, Higgs Session, Council Chamber

- **pseudo-scalar Higgs:**
NNLO cross section [R.H., Kilgore ('02)], [Anastasiou, Melnikov ('02)]
[Ravindran, Smith, v.Neerven ('03)]
NLO distributions [Field, Smith, Tejada-Yeomans, v.Neerven]


- **squark loops:** suppressed by $\frac{m_t^2}{\tilde{m}_t^2}$

NLO: [Dawson, Djouadi, Spira ('96)]




Supersymmetry: Bottom quarks


$$\frac{\lambda_b}{\lambda_t} = \frac{m_b}{m_t} \cdot \frac{v_u}{v_d} = \frac{m_b}{m_t} \cdot \tan \beta$$



NLO: [Spira, Djouadi, Graudenz, Zerwas ('95)]




→



+ higher orders


NLO:
[Maltoni, Sullivan,
Willenbrock ('03)]

at NNLO:
[R.H., Kilgore ('03)]



— collinear terms

$b\bar{b} \rightarrow H$ at NNLO



Backgrounds

- which tools available?

beyond LO: MCFM [Ellis, Campbell]

DIPHOX [Binoth, Giullet, Pilon, Werlen]

- where do we need higher orders?

- use NNLO signal with NLO background?

- Goal: collection/qualification of existing tools for BG's

→ talk by T. Binoth

Wed, 17.15, Higgs session, Council Chamber

Summary and Conclusions

- LHC challenges → scientific progress!
NNLO calculations, Monte Carlos, ...
- Standard Higgs:
all signal and most important background processes
under good control
- SUSY Higgs: on the way

Goals of Higgs Session:

- Specify open issues for signals/backgrounds
- Qualify existing and identify missing TOOLS
→ talks by E. Richter-Was, M. Spira, ...
Thu, 9.00am, Higgs Session, Council Chamber