

# *$t\bar{t}H/b\bar{b}H$ PRODUCTION @ LHC*

Michael Spira (PSI)

I  $t\bar{t}H$  Production

II  $b\bar{b}H$  Production

III Conclusions

Phys. Rev. Lett. 87 (2001) 201805 [hep-ph/0107081]


Nucl. Phys. B653 (2003) 151 [hep-ph/0211352]

in collaboration with W. Beenakker, S. Dittmaier,  
M. Krämer, B. Plümper and P.M. Zerwas

# I $t\bar{t}H$ PRODUCTION

- $gg, q\bar{q} \rightarrow t\bar{t}H$  important for  $M_H \lesssim 130$  GeV

F



- crucial for determination of top Yukawa coupling
- Equivalent Higgs Approximation:  $M_H^2 \ll m_t^2 \ll \hat{s}$

$$\hat{\sigma}(t\bar{t}H) = (f_{H/t} + f_{H/\bar{t}}) \otimes \hat{\sigma}(t\bar{t})$$



Dawson, Reina

valid within factor 2


QCD corrections  $\sim 20\%$  [scale =  $m_t + M_H/2!$ ]

- full calculation: two groups
- Beenakker, Dittmaier, Krämer, Plümper, S., Zerwas
- Dawson, Jackson, Orr, Reina, Wackerath

→ full agreement now for total cross sections




(i) Virtual Corrections





- most difficult part: Pentagon diagrams [infrared and collinear divergent]
- $\alpha_s$ :  $\overline{\text{MS}}$  scheme [5 flavours],  $m_t$ : on-shell



(ii) Real Corrections





- complex matrix elements
- infrared and collinear singularities cancel against virtual corrections and counter terms of PDFs [mass factorization]
- PDF:  $\overline{\text{MS}}$  scheme [5 flavours]



Beenakker, Dittmaier, Krämer, Plümper, S., Zerwas




Beenakker, Dittmaier, Krämer, Plümper, S., Zerwas



Beenakker, Dittmaier, Krämer, Plümper, S., Zerwas


# II $b\bar{b}H$ PRODUCTION

- $b\bar{b} + H/A$  dominant for large  $\tan\beta$




large logs from phase space integration  
 $\rightarrow$  absorbed in bottom PDF  
 resummation  $\equiv$  DGLAP evolution

- new processes:



NNLO



NLO



Dicus, Willenbrock  
 Stelzer, ...  
 Balazs, ...  
 Campbell, ...

Harlander, Kilgore

$$b(x, \mu^2) \longrightarrow b(x, \mu^2) - \frac{\alpha_s}{2\pi} P_{qg} \otimes g(x, \mu^2) \log \frac{\mu^2}{m_b^2}$$


$$\mu \sim M_\phi \Rightarrow \sigma_{tot}$$






Harlander, Kilgore



$p\bar{p} \rightarrow hb$  @ Tevatron



$p\bar{p} \rightarrow hb$  @ Tevatron



Campbell, Ellis, Maltoni, Willenbrock



- factorization in high-energy limit: [ $M_{Tb} = \sqrt{p_{Tb}^2 + m_b^2}$ ]

$$\frac{d\sigma^{(2\rightarrow 3)}}{dM_{Tb}^2} = \frac{1}{M_{Tb}^2} \left\{ \frac{\alpha_s}{2\pi} \Delta_{qg} \otimes g \otimes g \otimes \hat{\sigma}_{\bar{b}g} \right\}_{M_{Tb}=m_b \rightarrow 0} + \dots$$

$$\Delta_{qg}(x) = P_{qg}(x) + \frac{m_b^2}{M_{Tb}^2} x(1-x)$$

- total cross section:

$$\sigma = \underbrace{\int_{m_b^2}^{\mu_F^2} \frac{dM_{Tb}^2}{M_{Tb}^2}}_{\log \frac{\mu_F^2}{m_b^2}} \left\{ \frac{\alpha_s}{2\pi} P_{qg} \otimes g \otimes g \otimes \hat{\sigma}_{\bar{b}g} \right\}_{M_{Tb}=m_b \rightarrow 0} + \dots$$

$\Rightarrow$  crucial condition:



F

$$\frac{d\sigma^{(2\rightarrow 3)}}{dM_{Tb}} \propto \frac{1}{M_{Tb}} \quad \text{up to } M_{Tb} \sim \mu_F$$



- $p_T > p_{Tmin}$ :  $\log \frac{\mu_F^2}{m_b^2} \rightarrow \log \frac{\mu_F^2}{M_{Tmin}^2}$

$\Rightarrow$  unintegrated PDFs...



- large corrections to  $gg \rightarrow b\bar{b}H$ ?



Rainwater, S., Zeppenfeld



Dittmaier, Krämer, S. [preliminary]



Dittmaier, Krämer, S. [preliminary]  
 [compared to Maltoni, Sullivan, Willenbrock]

# III CONCLUSIONS

## (i) $t\bar{t}H$

- QCD corrections:  $\sim (20 - 40)\%$  @ LHC
- strong reduction of scale dependence:  
 $\Delta \lesssim 10 - 15\%$
- shape of Higgs  $p_T$  distribution changed by 10%  
[scale = transverse mass]
- shape of (anti)top  $p_T$  distribution hardly affected  
[scale = transverse mass]
- shape of rapidity distributions hardly modified

## (ii) $b\bar{b}H$

- QCD corrections: (50–120)% for total cxn  
 $\mathcal{O}(20-40)\%$  with 2  $b$  jets
- reasonable agreement with  $b\bar{b} \rightarrow H$  [ $\mu_F = M_H/4$ ]
- reliability of bottom PDFs still not clear [scale choice, approximate kinematics, etc.]  
 $\Rightarrow$  more [conceptual] work required