HERA-LHC Charm and Bottom

Matteo Cacciari, Massimo Corradi, Andrea Dainese, Andreas Meyer, Maria Smizanska, Ulrich Uwer, Christian Weiser

Experimental and theoretical overview

Measurements at HERA, open HF, quarkonium

Prospects for LHC

Triggers for HF physics

Topics to be studied

HF Triggers


Comparisons of strategies in different environments of HERA and LHC machines

Main complaint: simulation of the background is unsatisfactory. Large uncertainties for LHC

Interesting cross-talk of two communities.
Unclear if it will be pursued any further in the workshop

Minimum bias and underlying event for HF triggers Painful point for simulations. We hope other groups (MC? JETS?) will address this

Poster plot:



Yes, HERA can measure HQ production, and the theory can describe it

Topics to address during the Workshop

Small-x, k_T factorization?

Not discussed in this meeting for lack of time

Interesting subject for follow-up meeting (e.g. for bottom cross section at LHC)

Correlations (c-cbar and b-bbar)

Present tools: NLO codes, MC

MC@NLO strongly asked for (demanded?) by HERA experimental community

Experimental methods: what and how to measure at HERA and LHC

MC tuning for HF using HERA data + JetWEB and predictions for LHC



MC group + LCG

Heavy Quarkonium


J/psi and Upsilon

Review of HERA results

Not fully satisfying situation concerning comparisons with NRQCD theory and Tevatron data

Hopes for clarifying the situation before LHC starts?

Validation of QCD techniques for HQ PDFs / Resummation




Ingredient: photon PDFs improvements

Tests of fragmentation/validation in hadronic environment

HQ FF fitted at LEP and used at the Tevatron. Let's check at HERA too

Comparison of charm FF at ARGUS, ZEUS and OPAL



See you in DESY