

# ***Status and Prospects of Rare $K^\pm$ and $K_L$ Decays from NA48***

Rainer Wanke

Institut für Physik, Universität Mainz

**Kaon Mini Workshop**

CERN, May 5, 2004

# Introduction

1997	$\varepsilon'/\varepsilon$ run	$K_L + K_S$
1998	$\varepsilon'/\varepsilon$ run	$K_L + K_S$
1999	$\varepsilon'/\varepsilon$ run $K_L + K_S$	$K_S$ Hi. Int.
2000	$K_L$ only	$K_S$ High Intensity <i>NO Spectrometer</i>
2001	$\varepsilon'/\varepsilon$ run $K_L + K_S$	$K_S$ High Int.
2002	$K_S$ High Intensity	
2003	$K^\pm$ High Intensity	
2004	$K^\pm$ High Intensity	

## $K^\pm$ Decays: A Look into the Future

- **NA48/2** Data taking 2003 (+ 2004):  
World largest multi-purpose sample  
of  $K^\pm$  decays.

## $K_L$ Decays: Wrapping up the Past

- **NA48** Data taking up to 2001:  
Huge  $K_L$  sample, several publications,  
still some left-overs.

## Aim of this talk:

- Introduce analyses which are expected to be finished soon.
- Estimates of sample sizes, backgrounds, and expected uncertainties.
- But: Mostly no results yet.

## ■ $K^\pm$ Decays:


- $K^\pm \rightarrow \pi^\pm \pi^\mp e^\pm \nu (\bar{\nu})$  ( $K_{e4}^+$ ) branching ratio and form factors
- $K^\pm \rightarrow \pi^\pm e^+ e^-$  and  $K^\pm \rightarrow \pi^\pm \mu^+ \mu^-$  decays
- First look into  $K^\pm \rightarrow \pi^\pm \gamma\gamma$

## ■ $K_L$ Decays:

- $K_L \rightarrow \pi^\pm \pi^0 e^\mp \nu_e$  ( $K_{e4}^0$ ) branching ratio and form factors
- Form factor in  $K_L \rightarrow e^+ e^- \gamma$
- Measurement of  $K_L \rightarrow e^+ e^- e^+ e^-$

## ■ Not covered here:

- CP violation in  $K^\pm \rightarrow \pi^\pm \pi^\pm \pi^\mp$   $\implies$  Spasimir Balev
- $K_S$  and neutral hyperon decays  $\implies$  Cristina Lazzeroni
- Semileptonic decays  $K_{l3}, K_{l3\gamma}$   $\implies$  Mayda Velasco



## Main detector components:


- Magnetic spectrometer:  
 $\Delta p/p \simeq 0.5\% \oplus 0.009\% \times p [\text{GeV}/c]$
- Liquid-krypton el.-magn. calorimeter  
 $\Delta E/E \simeq 3.2\%/\sqrt{E} \oplus 9\%/E \oplus 0.42\%$
- Hadron calorimeter, photon veto,  $\mu$  veto counters

# Rare $K^\pm$ Decays

# NA48/2 $K^\pm$ Data Taking in 2003

## NA48/2 experiment in 2003:

- High intensity data taking with simultaneous  $K^+$  and  $K^-$  beams.
- New beam spectrometer (Micromesh gas chambers)  
     $\Rightarrow K^\pm$  momentum resolution  $\sim 1\%$
- $\sim 50$  days of data taking:  $\sim 10^{11} K^\pm$  decays in decay volume.



## Trigger:

- All 3-track events  
( $\sim 98\%$  efficient)
- 1-track events  
with  $\mu$ -veto and  
$$(p_K - p_\pi)^2 \gg m_{\pi^0}^2$$
  
to reject  $K^\pm \rightarrow \pi^\pm \pi^0$ .

# NA48/2 $K^\pm$ Data Taking in 2003

- Total data taking period 2003:  $\sim 50$  days.
- Data presented here: Last month of data taking  
(so-called Super Samples 1,2,3).
  - ➡ Stable running conditions during this period.
- All 2003 data almost twice as much.
  - ➡ Still useful for Rare Decay analyses.

# **Measurement of $K^+ \rightarrow \pi^+\pi^-e^+\nu$ ( $K_{e4}^+$ )**

## ■ Physics interest:

- Low energy  $\pi\pi$  scattering unambiguously predicted from Chiral Perturbation Theory first principles.
- *s*-wave isospin zero  $\pi\pi$  scattering length  $a_0^0$  can be determined from form factors in  $K_{e4}^+$  decays.
  - ⇒ Predicted to  $a_0^0 = 0.220 \pm 0.005$  in ChPT.  
(Colangelo, Gasser, Leutwyler, 2001)
  - ⇒ Direct and firm test of Chiral Perturbation Theory.


## ■ Previous measurements:

- Geneva-Saclay (1977): 30 000 events
- Brookhaven E865 (2001): 400 000 events
  - ⇒  $a_0^0 = 0.216 \pm 0.013$ .

# Measurement of $K^+ \rightarrow \pi^+\pi^-e^+\nu$ ( $K_{e4}^+$ )







- Events fully reconstructed by using known kaon momenta.  
➡ More than 300 000 events in SS 1–3.
- Background mainly from  $K^+ \rightarrow \pi^+\pi^+\pi^-$  and  $K^+ \rightarrow \pi^+\pi^0_{\text{Dalitz}}$  with  $\pi \leftrightarrow e$  mis-identification.  
➡ Use of neural network for  $e/\pi$  separation.
- Total background: 0.6%  
Determined from wrong-sign  $\pi^+\pi^+e^-$  data events.

Invariant kaon mass Ke4 hypothesis



# **Measurement of $K^+ \rightarrow \pi^+\pi^-e^+\nu$ ( $K_{e4}^+$ )**

**Cabibbo-Maksymowicz variables:** (Background in red)



# ***Measurement of $K^+ \rightarrow \pi^+\pi^-e^+\nu$ ( $K_{e4}^+$ )***

## **Status and Prospects for $K_{e4}^+$ :**

- Expect more than half a million events in total 2003 data.
- Backgrounds are tiny, systematics should not be dominating.
- Analysis underway, but a lot of work.
  - ➡ May have to wait til the end of the year.

# **Measurement of and $K^\pm \rightarrow \pi^\pm \mu^+ \mu^-$**



## ■ Physics interest:

- FCNC, suppressed by GIM mechanism.
- Amplitude dominated by long-distance contributions (one-photon “bremsstrahlung” diagrams).  
    ⇒ Can be extracted from form factor measurements!
- Potentially interesting channels for CP violation between  $\Gamma(K_{\pi ll}^+)$  and  $\Gamma(K_{\pi ll}^-)$ .

## ■ Previous measurements:

- $K^\pm \rightarrow \pi^\pm e^+ e^-$ : BNL E865 found 10500 events  
    ⇒  $\text{Br} = (2.94 \pm 0.05 \pm 0.14) \times 10^{-7}$   
(Also form factor measurement.)
- $K^\pm \rightarrow \pi^\pm \mu^+ \mu^-$ : Several experiments, in total 800 events.  
    ⇒  $\text{Br} = (0.81 \pm 0.14) \times 10^{-7}$


# Measurement of $K^\pm \rightarrow \pi^\pm e^+ e^-$



- About 2600 candidates for  $m_{ee} > 140 \text{ MeV}/c^2$  in one month of data taking.
- Background very small ( $\sim 1 - 2 \%$ ).

# ***Measurement of $K^\pm \rightarrow \pi^\pm e^+ e^-$***


## ■ **Form Factor:**



## ■ **Status and Prospects:**

- Statistical error on branching fraction  $\Delta\text{Br} \leq \pm 0.06 \times 10^{-7}$ .  
(Total PDG error now is  $\pm 0.13 \times 10^{-7}$ )
- Systematics expected to be small.
- Analysis far advanced.  
→ Expect result within the next months.

# ***Measurement of $K^\pm \rightarrow \pi^\pm \mu^+ \mu^-$***



- Event sample of  $> 10^3 K^\pm \rightarrow \pi^\pm \mu^+ \mu^-$  candidates (SS 1–3).  
(Current world sample: 800 events)
- Again: Very small background.

# Measurement of $K^\pm \rightarrow \pi^\pm \gamma\gamma$

- $K^\pm \rightarrow \pi^\pm \gamma\gamma$ :

Similar situation as for  $K_L \rightarrow \pi^0 \gamma\gamma$ :


- $\mathcal{O}(p^2)$  ChPT amplitude vanishes.
- $\mathcal{O}(p^4)$  computed  
(but not exactly known for  $K^\pm \rightarrow \pi^\pm \gamma\gamma$ !)
- Signal mainly has  
 $m_{\gamma\gamma} > 2m_\pi$
- Fit to  $m_{\gamma\gamma}$  distribution:  
    → Information on  $\mathcal{O}(p^6)$

- **Experimental problem:**

Suppression of  $K^\pm \rightarrow \pi^\pm \pi^0$  at trigger level.

    → Two trigger conditions for  $K^\pm \rightarrow \pi^\pm \gamma\gamma$  events:


- Normal neutral trigger → down-scaled by 80.
- “High- $m_{\gamma\gamma}$ ” trigger → no down-scaling, but no low- $m_{\gamma\gamma}$






# *Measurement of $K^\pm \rightarrow \pi^\pm \gamma\gamma$*

**First look into the data:** (Compare with 31 events of E787!)

**Neutral trigger (DS=80):**



**High- $M(\text{gg})$  trigger (DS=1):**



➡ Promising — but still a lot of work to do!

# Rare $K_L$ Decays

# Rare $K_L$ Decays from NA48

- Main goal of **NA48/0**:

Measurement of  $\text{Re}(\epsilon'/\epsilon) = (14.7 \pm 2.2) \times 10^{-4}$ .

⇒ Data taking finished in 2001, published in 2002.

- **Meanwhile:** Several rare decay analyses published.

(E.g.  $K_L \rightarrow \pi^0 \gamma \gamma$ ,  $K_{L,S} \rightarrow \pi^+ \pi^- e^+ e^-$ , rare  $K_S$  decays, ...)

- Still some **left-overs** due to **complicated analyses** and **available man-power**.

⇒ Now being finished!

- $K_L \rightarrow \pi^\pm \pi^0 e^\mp \nu_e$  ( $K_{e4}^0$ )

- $K_L \rightarrow e^+ e^- \gamma$

- $K_L \rightarrow e^+ e^- e^+ e^-$







- $K_{e3}, K_{\mu 3}$  ⇒ Mayda Velasco

# **Measurement of $K_L \rightarrow \pi^\pm \pi^0 e^\mp \nu_e$ ( $K_{e4}^0$ )**

- **Physics interest:** Similar, but not identical, to  $K_{e4}^+$ .
  - Measurement of branching fraction and form factors allow determination of ChPT parameters as e.g. the coefficient  $L_3$ .
- **Previous measurements:**
  - E731 (Fermilab, 1993): 729 events  
(Measured both Br and form factors.)
- **NA48 2001:**
  - Two different triggers:
    - Neutral minimum-bias (downscaled by 30)
    - Special  $K_{e4}$  trigger (downscaled by 50)
  - Total  $K_{e4}$  sample: 5464 events (bkg contamination only 1.1%)
    - ⇒ One order of magnitude more precise than previous measurement!

# Measurement of $K_L \rightarrow \pi^\pm \pi^0 e^\mp \nu_e$ ( $K_{e4}^0$ )

## Fit of Cabibbo-Maksymowicz variables:



# **Measurement of $K_L \rightarrow \pi^\pm \pi^0 e^\mp \nu_e$ ( $K_{e4}^0$ )**

## **Results on $K_{e4}^0$ :** (submitted to PLB)

- Branching fraction:

$$\text{Br}(K_{e4}^0) = (5.21 \pm 0.07_{\text{stat.}} \pm 0.09_{\text{syst.}}) \times 10^{-5}$$

- Form factors:

$$\overline{f}_s = 0.052 \pm 0.006_{\text{stat.}} \pm 0.002_{\text{syst.}}$$

$$\overline{f}_p = -0.051 \pm 0.011_{\text{stat.}} \pm 0.005_{\text{syst.}}$$

$$\lambda_g = 0.087 \pm 0.019_{\text{stat.}} \pm 0.006_{\text{syst.}}$$

$$\overline{h} = -0.32 \pm 0.12_{\text{stat.}} \pm 0.05_{\text{syst.}}$$


in good agreement with ChPT predictions.

- Determination of ChPT coefficient  $L_3$  (from branching ratio):



$$L_3 = (-4.1 \pm 0.2) \times 10^{-3}$$

# **Form Factor in $K_L \rightarrow e^+e^-\gamma$**

- NA48 publication (2000) on 97 data:  
BMS form factor:  $\alpha_{K^*} = 0.36 \pm 0.06$ .
- Result in slight contradiction with KTeV measurements  
on  $K_L \rightarrow e^+e^-\gamma$  and  $K_L \rightarrow \mu^+\mu^-\gamma$   
on a much larger data sample.
- Now: Using total NA48 data set  
(97,98,99,2001)  
⇒ About a factor of 9  
statistical improvement  
w.r.t. 97 publication.




# Form Factor in $K_L \rightarrow e^+ e^- \gamma$



- $\approx 60\,000$  events in total NA48 data (Background  $\sim 1\%$ )
  - Statistical error:  $\Delta\alpha_{K^*} \approx 0.02$
- Huge amount of work was done on radiative corrections.
- Expect result for BMS and DIP models within the next months.

# ***Measurement of $K_L \rightarrow e^+e^-e^+e^-$***

- $K_L \rightarrow e^+e^-e^+e^-$  measurement soon to be finished.
- 200 events found with expected background of 1%.



# Conclusions

## ■ Rare $K^\pm$ decays:

All measurements with large statistics and low systematics.

Lots of on-going work:

- High precision measurement of  $K_{e4}^+$ .
- Branching fraction and form factors of  $K^\pm \rightarrow \pi^\pm e^+ e^-$  and  $K^\pm \rightarrow \pi^\pm \mu^+ \mu^-$ .
- Work on  $K^+ \rightarrow \pi^+ \gamma\gamma$ .
- Many more analyses not shown here:

$K_{\mu 4}$ ,  $K^\pm \rightarrow \pi^\pm \pi^0 \gamma$ ,  $K^\pm \rightarrow \pi^\pm \gamma e^+ e^-$ ,  $K^\pm \rightarrow l^\pm l^+ l^- \nu$ , ...

## ■ $K_L$ decays:

- Precision measurement on  $K_{e4}^0$  finished.
- Measurements of  $K_L \rightarrow e^+ e^- \gamma$  and  $K_L \rightarrow e^+ e^- e^+ e^-$  on whole data set expected to be published in summer.