


# NA49-future **Astroparticle physics programme**



**From Quarks to Cosmos** 

still an open question:


cosmic ray flux spectrum

- sources
- acceleration
- transport
- highest energies

Andreas Haungs Institut für Kernphysik Forschungszentrum Karlsruhe Karlsruhe, Germany haungs@ik.fzk.de



## Astroparticle physics programme Air shower measurements




No direct measurements for  $E_0 > 10^{14} \text{ eV}$ 

→ Air shower measurements


Air shower development:

- → longitudinal distribution
- → lateral distribution






## Astroparticle physics programme Shower observables







## Astroparticle physics programme Problem of the knee






## Astroparticle physics programme Muon distribution








## Astroparticle physics programme Muon distribution






## Astroparticle physics programme Problem of highest energy






## Astroparticle physics programme Highest energy event



highest energy event measured by the Pierre Auger Experiment


#### **Energy of cosmic rays**

- -Signal density lateral distribution
- -muon number dominated






## Astroparticle physics programme Muon lateral distribution




Energy of cosmic rays -muon lateral density



muon lateral density distributions as predicted by hadronic interaction models



## Astroparticle physics programme Reactions to be measured



#### NA49-future:

Pion spectra and multiplicities in

p-Air

π-Air

p-p

p-C

C-C

at SPS energies

needed for understanding the whole cosmic ray flux spectrum

default transition energy of low and high energy interaction models in CORSIKA

pion multiplicity for p-p interactions (Eichten et al.) compared to simulations