



Search for Randall-Sundrum Gravitons in CMS

C. Collard (LLR, Ecole Polytechnique)
and M.-C. Lemaire (Dapnia, Saclay)

The Randall-Sundrum Model

The e^+e^- Analysis

Correction for the electronics saturation

Search for massive resonances

Results & Conclusions



The Randall-Sundrum Model



One Warped Extra Dimension = Answer to the Hierarchy Problem

- 5D Anti-de-Sitter space-time with 2 branes of 4D:

Metric: $e^{-2kr\varphi} \eta_{\mu\nu} dx^\mu dx^\nu + r^2 d\varphi^2$

Curvature: $k (\sim M_{PL})$


Compactification radius: r

New coordinate: $\varphi (-\pi \leq \varphi \leq \pi)$

Traditional 4D coordinates: x^μ

- Gravity scale : $\Lambda_\pi = M_{PL} e^{-kr\pi}$

no new hierarchy with $\Lambda_\pi \sim 1 \text{ TeV}$
if $kr \approx 11-12$





The Randall-Sundrum Model




Only the graviton can propagate in 5D. On the 4D branes, Kaluza-Klein excitations of the graviton can be observed:

$$M_n = k x_n e^{-kr\pi} \text{ avec } J_1(x_n)=0$$

$$\Gamma_n = \rho M_n x_n^2 c^2$$

with two free parameters in the model: $M_G = M_1$ and $c = k/M_{PL}$

Example: if $M_1 = 1.5 \text{ TeV}$



H.Davoudiasl, J.Hewett,
T.Rizzo, hep-ph/0006041

- $c=1.$
- $c=0.5$
- $c=0.1$
- $c=0.05$
- $c=0.01$


Look for the 1st KK graviton (resonance @ $M_G \sim \text{TeV}$)



The Randall-Sundrum Model



Constraints on the two free parameters of the model: M_G and $c=k/M_{Pl}$



H.Davoudiasl, J.Hewett,
T.Rizzo, hep-ph/0006041

Which part of the plane can be access with CMS?




The e^+e^- channel



- **Signal:** $pp \rightarrow G \rightarrow e^+e^-$ (K Factor =1)

The e^+e^- decay channel has a low branching ratio (BR=2%) but the clear signal in the electromagnetic calorimeter ECAL allows it to be the **discovery channel for Randall-Sundrum Gravitons**.



- **Background:** 2 electrons in the final state
 - Drell-Yan: $pp \rightarrow \gamma/Z \rightarrow e^+ e^-$ (K Factor=1.3)
 - [Jet faking an electron: Dijet, γ -jet, e -jet which is negligible in comparison to Drell-Yan]



Technical details



- Generation with PYTHIA (+ inner Bremsstrahlung with PHOTOS)
- **Full Simulation and Reconstruction chain of CMS**
(CMSIM & ORCA without pile-up):
 - Synchrotron radiation is included but found to be negligible in comparison to Bremsstrahlung in the tracker
 - Work on the electron reconstruction
 - Possible saturation of the ECAL electronics (pre-amplifiers in VFE cards) is studied:
 - Saturation expected at 1.7 TeV in the barrel with measured crystal light yield (4.5 photo-electrons/MeV)
 - Study here for saturation at 1.25 TeV (i.e. 6 p.e./MeV)
 - A simple correction is found.






Saturation of the ECAL electronics



- The saturation has a big effect on the mass reconstruction of heavy resonances.

Idea for correction:
Correlation between
 $Red_5 = E_9 - E_4$ and E_1

5x5 crystals






Saturation of the ECAL electronics




- This correction of the saturation allows to reconstruct heavy mass resonances.





Selection Cuts





$$pp \rightarrow G \rightarrow e^+e^-$$

- Trigger up to Level 2.5
- 2 electrons
 - Super-Clusters:
 - $p_T > 100 \text{ GeV}$,
 - $|\eta| < 1.4442$ (barrel)
or $1.566 < |\eta| < 2.5$ (endcaps)
 - Isolated: $E_T^{\text{cone}} < 0.02 E_T^{\text{SC}}$ in cone $\Delta r < 0.5$ (to kill big jets)
 - Electromagnetic: $H/E < 0.1$ (to kill π^+/π^-)
 - Charged: 2 tracks with at least 2 hits (to kill π^0/γ)




Search for a resonance



- Fit of a Gaussian to the signal distribution
- Mass window for N_S and N_B estimation: $\langle M \rangle \pm 3\sigma$
- For low coupling values: $E_1 < 1.25 \text{ TeV}$ (no saturation)
- For large coupling values: correction of the saturation coming from the ECAL electronics




Results for $c=0.01$





Results for $c=0.01$







Significance




$$S = 2(\sqrt{N_S + N_B} - \sqrt{N_B}).$$





$G \rightarrow e^+e^-$: Discovery plane





Conclusions



Full simulation & reconstruction analysis

- Study of very energetic electrons and search for massive resonances
- Discovery plane for the Randall-Sundrum gravitons $G \rightarrow e^+ e^-$:
 - With 100 fb^{-1} : the region of interest will be covered by CMS.
 - With 1 fb^{-1} : a large part of this region of interest will be accessible at the first beginning of the LHC running.
- For the Future: Work on the Identification of the Graviton nature
 - Angular Distribution (Graviton is spin 2)
 - Other channels:
 - $G \rightarrow \gamma\gamma$ is allowed but not $Z' \rightarrow \gamma\gamma$.
 - Test the universality of the Graviton couplings.