CP Violation in Charged Higgs Production

Jennifer Williams

Cavendish Laboratory, University of Cambridge

2 December 2004

Structure ...

- Introduction
- Decay
- Production
- Work in Progress

CP violation in charged Higgs production?

- CP violation in charged Higgs production?
- Origin is in soft susy breaking terms...

- CP violation in charged Higgs production?
- Origin is in soft susy breaking terms...

$$\begin{aligned} -\mathcal{L}_{soft} &= \frac{1}{2} \left(M_{3} \tilde{g} \tilde{g} + M_{2} \tilde{W} \tilde{W} + M_{1} \tilde{B} \tilde{B} + \text{h.c.} \right) \\ &+ \mathbf{m}_{\tilde{\mathbf{Q}}}^{2} \tilde{Q}^{\dagger} \tilde{Q} + \mathbf{m}_{\tilde{\mathbf{u}}}^{2} \tilde{u}^{c^{*}} \tilde{u}^{c} + \mathbf{m}_{\tilde{\mathbf{d}}}^{2} \tilde{d}^{c^{*}} \tilde{d}^{c} + \mathbf{m}_{\tilde{\mathbf{L}}}^{2} \tilde{L}^{\dagger} \tilde{L} + \mathbf{m}_{\tilde{\mathbf{e}}}^{2} \tilde{e}^{c^{*}} \tilde{e}^{c} \\ &+ \mathbf{A}_{\mathbf{u}} \tilde{u}^{c} \tilde{Q} H_{u} - \mathbf{A}_{\mathbf{d}} \tilde{d}^{c} \tilde{Q} H_{d} - \mathbf{A}_{\mathbf{e}} \tilde{e}^{c} \tilde{L} H_{d} + \text{h.c.} \\ &+ m_{H_{u}}^{2} H_{u}^{*} H_{u} + m_{H_{d}}^{2} H_{d}^{*} H_{d} + (b H_{u} H_{d} + \text{h.c.}) \end{aligned}$$

- CP violation in charged Higgs production?
- Origin is in soft susy breaking terms...

$$\begin{aligned} -\mathcal{L}_{soft} &= \frac{1}{2} \left(M_{3} \tilde{g} \tilde{g} + M_{2} \tilde{W} \tilde{W} + M_{1} \tilde{B} \tilde{B} + \text{h.c.} \right) \\ &+ \mathbf{m}_{\tilde{\mathbf{Q}}}^{2} \tilde{Q}^{\dagger} \tilde{Q} + \mathbf{m}_{\tilde{\mathbf{u}}}^{2} \tilde{u}^{c^{*}} \tilde{u}^{c} + \mathbf{m}_{\tilde{\mathbf{d}}}^{2} \tilde{d}^{c^{*}} \tilde{d}^{c} + \mathbf{m}_{\tilde{\mathbf{L}}}^{2} \tilde{L}^{\dagger} \tilde{L} + \mathbf{m}_{\tilde{\mathbf{e}}}^{2} \tilde{e}^{c^{*}} \tilde{e}^{c} \\ &+ \mathbf{A}_{\mathbf{u}} \tilde{u}^{c} \tilde{Q} H_{u} - \mathbf{A}_{\mathbf{d}} \tilde{d}^{c} \tilde{Q} H_{d} - \mathbf{A}_{\mathbf{e}} \tilde{e}^{c} \tilde{L} H_{d} + \text{h.c.} \\ &+ m_{H_{u}}^{2} H_{u}^{*} H_{u} + m_{H_{d}}^{2} H_{d}^{*} H_{d} + (b H_{u} H_{d} + \text{h.c.}) \end{aligned}$$

Need non trivial complex phases.

- CP violation in charged Higgs production?
- Origin is in soft susy breaking terms...

$$\begin{aligned} -\mathcal{L}_{soft} &= \frac{1}{2} \left(M_{3} \tilde{g} \tilde{g} + M_{2} \tilde{W} \tilde{W} + M_{1} \tilde{B} \tilde{B} + \text{h.c.} \right) \\ &+ \mathbf{m}_{\tilde{\mathbf{Q}}}^{2} \tilde{Q}^{\dagger} \tilde{Q} + \mathbf{m}_{\tilde{\mathbf{u}}}^{2} \tilde{u}^{c^{*}} \tilde{u}^{c} + \mathbf{m}_{\tilde{\mathbf{d}}}^{2} \tilde{d}^{c^{*}} \tilde{d}^{c} + \mathbf{m}_{\tilde{\mathbf{L}}}^{2} \tilde{L}^{\dagger} \tilde{L} + \mathbf{m}_{\tilde{\mathbf{e}}}^{2} \tilde{e}^{c^{*}} \tilde{e}^{c} \\ &+ \mathbf{A}_{\mathbf{u}} \tilde{u}^{c} \tilde{Q} H_{u} - \mathbf{A}_{\mathbf{d}} \tilde{d}^{c} \tilde{Q} H_{d} - \mathbf{A}_{\mathbf{e}} \tilde{e}^{c} \tilde{L} H_{d} + \text{h.c.} \\ &+ m_{H_{u}}^{2} H_{u}^{*} H_{u} + m_{H_{d}}^{2} H_{d}^{*} H_{d} + (b H_{u} H_{d} + \text{h.c.}) \end{aligned}$$

- Need non trivial complex phases.
- **Possible complex terms are** M_3 , μ and **A**.

- CP violation in charged Higgs production?
- Origin is in soft susy breaking terms...

$$\begin{aligned} -\mathcal{L}_{soft} &= \frac{1}{2} \left(M_{3} \tilde{g} \tilde{g} + M_{2} \tilde{W} \tilde{W} + M_{1} \tilde{B} \tilde{B} + \text{h.c.} \right) \\ &+ \mathbf{m}_{\tilde{\mathbf{Q}}}^{2} \tilde{Q}^{\dagger} \tilde{Q} + \mathbf{m}_{\tilde{\mathbf{u}}}^{2} \tilde{u}^{c^{*}} \tilde{u}^{c} + \mathbf{m}_{\tilde{\mathbf{d}}}^{2} \tilde{d}^{c^{*}} \tilde{d}^{c} + \mathbf{m}_{\tilde{\mathbf{L}}}^{2} \tilde{L}^{\dagger} \tilde{L} + \mathbf{m}_{\tilde{\mathbf{e}}}^{2} \tilde{e}^{c^{*}} \tilde{e}^{c} \\ &+ \mathbf{A}_{\mathbf{u}} \tilde{u}^{c} \tilde{Q} H_{u} - \mathbf{A}_{\mathbf{d}} \tilde{d}^{c} \tilde{Q} H_{d} - \mathbf{A}_{\mathbf{e}} \tilde{e}^{c} \tilde{L} H_{d} + \text{h.c.} \\ &+ m_{H_{u}}^{2} H_{u}^{*} H_{u} + m_{H_{d}}^{2} H_{d}^{*} H_{d} + (b H_{u} H_{d} + \text{h.c.}) \end{aligned}$$

- Need non trivial complex phases.
- Possible complex terms are M_3 , μ and A.
- Choose to investigate A_t , keeping other phases zero.





• CP violation in loop correction to tbH^{\pm} vertex.



Decay done by Christova, Eberl, Kraml and Majerotto (Nucl.Phys. B639 (2002) 263-280; Erratum-ibid. B647 (2002) 359-360)



- Decay done by Christova, Eberl, Kraml and Majerotto (Nucl.Phys. B639 (2002) 263-280; Erratum-ibid. B647 (2002) 359-360)
- Decided to look at production



- Decay done by Christova, Eberl, Kraml and Majerotto (Nucl.Phys. B639 (2002) 263-280; Erratum-ibid. B647 (2002) 359-360)
- Decided to look at production
- Aim: combine production and decay



- Decay done by Christova, Eberl, Kraml and Majerotto (Nucl.Phys. B639 (2002) 263-280; Erratum-ibid. B647 (2002) 359-360)
- Decided to look at production
- Aim: combine production and decay
- Used SPS pt 1a for initial study



- Decay done by Christova, Eberl, Kraml and Majerotto (Nucl.Phys. B639 (2002) 263-280; Erratum-ibid. B647 (2002) 359-360)
- Decided to look at production
- Aim: combine production and decay
- Used SPS pt 1a for initial study
- Calculate CP asymmetry:

$$\mathcal{A}_{CP} = \frac{\sigma(H^+) - \sigma(H^-)}{\sigma(H^+) + \sigma(H^-)}$$

To check my methods, computer code etc, did the decay.

- To check my methods, computer code etc, did the decay.
- Agreed for subdominant loops, eg $\tilde{t}\tilde{b}\tilde{\chi}^0$ and $\tilde{\chi}^0\tilde{\chi}^\pm\tilde{t}/\tilde{b}$

- To check my methods, computer code etc, did the decay.
- Agreed for subdominant loops, eg $\tilde{t}\tilde{b}\tilde{\chi}^0$ and $\tilde{\chi}^0\tilde{\chi}^{\pm}\tilde{t}/\tilde{b}$
- Discrepancy in dominant loop, due to conjugation error in other study.



- To check my methods, computer code etc, did the decay.
- Agreed for subdominant loops, eg $\tilde{t}\tilde{b}\tilde{\chi}^0$ and $\tilde{\chi}^0\tilde{\chi}^{\pm}\tilde{t}/\tilde{b}$



- To check my methods, computer code etc, did the decay.
- Agreed for subdominant loops, eg $\tilde{t}\tilde{b}\tilde{\chi}^0$ and $\tilde{\chi}^0\tilde{\chi}^{\pm}\tilde{t}/\tilde{b}$



Result is less asymmetry than in other study.

Having checked decay, did production.

- Having checked decay, did production.
- Two main processes for the LHC —

- Having checked decay, did production.
- Two main processes for the LHC Bottom – gluon fusion: $\overline{b}_{g}, \overline{b}_{g}, \overline{b}, \overline{b}_{g}, \overline{b}, \overline{b}_{g}, \overline{b}, \overline{b}, \overline{$

 \boldsymbol{Q}

- Having checked decay, did production.
- Two main processes for the LHC Bottom – gluon fusion: $\overline{b}_{g}, q, \overline{b}$ $\overline{b}_{g}, q, \overline{b}$ \overline{t} \overline{t}

Gluon – gluon fusion:



- Having checked decay, did production.
- Two main processes for the LHC Bottom – gluon $\overline{b} = \overline{b} = \overline{b}$ \overline{t} \overline{t} $\overline{b} = \overline{t}$

Gluon – gluon fusion:



Dominant process is bottom – gluon fusion.

- Having checked decay, did production.
- Two main processes for the LHC Bottom – gluon \overline{b} \overline{b} \overline{t} \overline{t} \overline{b} t \overline{t}

Gluon – gluon fusion:



- Dominant process is bottom gluon fusion.
- Need to take care when combining to avoid double counting.

 Worked out cross sections using FormCalc, by Thomas Hahn

Comput. Phys. Commun. **118** (1999) 153

Worked out cross sections using FormCalc, by Thomas Hahn

Comput. Phys. Commun. 118 (1999) 153

For parton level, get similar thresholds to decay case



Worked out cross sections using FormCalc, by Thomas Hahn

Comput. Phys. Commun. 118 (1999) 153

For parton level, get similar thresholds to decay case



• Production < decay and $\tilde{t}\tilde{b}\tilde{\chi}^0$ loop < $\tilde{t}\tilde{b}\tilde{g}$ loop

Need to include pdfs to get hadronic cross section.

- Need to include pdfs to get hadronic cross section.
- Did for varying charged Higgs mass to match decay.



- Need to include pdfs to get hadronic cross section.
- Did for varying charged Higgs mass to match decay.



Discrepancies when compared with D-W. Jung, K. Y. Lee and H. S. Song. hep-ph/0307246

- Need to include pdfs to get hadronic cross section.
- Did for varying charged Higgs mass to match decay.



- Discrepancies when compared with D-W. Jung, K. Y. Lee and H. S. Song. hep-ph/0307246
- These are under investigation!

- Need to include pdfs to get hadronic cross section.
- Did for varying charged Higgs mass to match decay.



- Discrepancies when compared with D-W. Jung, K. Y. Lee and H. S. Song. hep-ph/0307246
- These are under investigation!
- Very small asymmetry challenging to see at LHC.

• Cross section $\sim 0.050 \text{ pb}$ for low Higgs mass $\sim 400 \text{ GeV}$.

- Cross section $\sim 0.050 \text{ pb}$ for low Higgs mass $\sim 400 \text{ GeV}$.
- Consistent with result from HERWIG.

- Cross section $\sim 0.050 \text{ pb}$ for low Higgs mass $\sim 400 \text{ GeV}$.
- Consistent with result from HERWIG.
- For integrated luminosity of $100 \text{ fb}^{-1} \rightarrow 5000 \text{ events.}$

- Cross section $\sim 0.050 \text{ pb}$ for low Higgs mass $\sim 400 \text{ GeV}$.
- Consistent with result from HERWIG.
- For integrated luminosity of $100 \text{ fb}^{-1} \rightarrow 5000 \text{ events.}$
- Much reduced by acceptance level, say $5\% \rightarrow 250$ events.

- Cross section $\sim 0.050 \text{ pb}$ for low Higgs mass $\sim 400 \text{ GeV}$.
- Consistent with result from HERWIG.
- For integrated luminosity of $100 \text{ fb}^{-1} \rightarrow 5000 \text{ events.}$
- Much reduced by acceptance level, say $5\% \rightarrow 250$ events.
- Not looking promising! Probably need to wait for luminosity upgrade.

Include gluon – gluon fusion.

- Include gluon gluon fusion.
- Explore other parts of parameter space:

- Include gluon gluon fusion.
- Explore other parts of parameter space:
 - Different SPS points 1b partially completed

- Include gluon gluon fusion.
- Explore other parts of parameter space:
 - Different SPS points 1b partially completed
 - Vary different parameters ϕ_{A_t} partially completed

- Include gluon gluon fusion.
- Explore other parts of parameter space:
 - Different SPS points 1b partially completed
 - Vary different parameters ϕ_{A_t} partially completed
 - Investigate other loops
 e.g., self energy loops, box diagram



- Include gluon gluon fusion.
- Explore other parts of parameter space:
 - Different SPS points 1b partially completed
 - Vary different parameters ϕ_{A_t} partially completed
 - Investigate other loops
 e.g., self energy loops, box diagram



Explore possibilities at International Linear Collider cleaner environment