

Analysis of the decay $B^\pm \rightarrow K^\pm \pi^\mp \pi^\pm$
at the *BABAR* Experiment

IOP Particle Physics 2004
6th April 2004

Thomas Latham
University of Bristol
BABAR Collaboration

Introduction

- Present results from the completed analysis:
 - Measurements of the **exclusive** branching fractions of B^\pm decays to $K^\pm \pi^\mp \pi^\pm$ final states
- Introduce current work:
 - **Amplitude analysis** of B^\pm decays to the final state $K^\pm \pi^\mp \pi^\pm$

Physics Motivation

- B^\pm decays to the final state $K^\pm \pi^\mp \pi^\pm$ via intermediate resonances can be used to search for **direct CP violation**
- Measurements of the branching fractions of the intermediate resonances can be compared with predictions from **hadronic models** (QCD Factorisation etc.), e.g.
 - W. N. Cottingham, *et al.*, J. Phys. **G28** (2002) 2843
 - M. Beneke and M. Neubert, Nucl. Phys. **B675** (2003) 333-415
 - C. Chiang *et al.*, Phys. Rev. D **69** (2004) 034001
 - R. Aleksan, *et al.*, Phys. Rev. D **67** (2003) 094019
- Study of these decays can also help to clarify the **nature of the resonances** involved, not all of which are well understood

General Analysis Issues 1 - Kinematic Variables

- $e^+e^- \rightarrow \Upsilon(4S) \rightarrow B\bar{B}$
 - B produced almost **at rest** in $\Upsilon(4S)$ frame
- Use **beam energy** to improve resolution
- Energy and momentum conservation give:
 - $\Delta E = E_B^* - E_{beam}^* \rightarrow 0$ for signal
 - $m_{ES} = \sqrt{E_{beam}^{*2} - p_B^{*2}} \rightarrow m_B$ for signal


General Analysis Issues 2 - Event Topology

- B produced almost **at rest** in $\Upsilon(4S)$ frame
 - B decays are **isotropic**
- $q\bar{q}$ decay products can have considerable momentum
 - Continuum (udsc) decays are **jet-like**
- Form a **Fisher Discriminant** of topological variables
 - This is a **linear combination** of variables
 - Gives **greater separation** between hypotheses than selecting on the variables alone

$B^\pm \rightarrow K^\pm \pi^\mp \pi^\pm$ Exclusive Branching Fraction Overview


- Investigate resonance composition
- Divide Dalitz Plot into regions
- Measure yields to each region
- Maximum Likelihood analysis with PDFs for m_{ES} , ΔE , Fisher
Parameterised separately for each region
- Then interpret yields in regions as BFs of resonances using
coupled resonance model considering interferences as a systematic
- Dataset: 61 million $B\bar{B}$ pairs

Regions of the Dalitz Plot




Fit Variable Projection Plots

- Histograms shown are for Region I (K^{*0})
- Histograms have likelihood ratio cut on other two variables
- Fit projections: **total** **background** **signal**




Dalitz Plots

- Plots have likelihood ratio cut in ΔE & Fisher
- Left-hand plot has **signal-like** cut in m_{ES}
- Right-hand plot has **background-like** cut in m_{ES}




Mass Projections

- Background subtracted projections of $m_{K\pi}$ & $m_{\pi\pi}$



Mass & Helicity Angle Projection Plots



- Histograms have likelihood ratio cut
- Have been background subtracted & efficiency corrected



Upper plots - Region I (K^{*0}); Lower Plots - Region V ($f_0(980)$)

The “higher K^{*0} ” region

- Structure cannot be explained by any single resonance
- Up to $m_{K\pi} = 1.6 \text{ GeV}/c^2$ shows similar structure to that observed on LASS for $K_0^{*0}(1430)$
 - D. Aston *et al.*, Nucl. Phys. **B296** (1988) 493



Branching Fractions

Channel	BF ($\times 10^{-6}$)	Errors ($\times 10^{-6}$)			
		Stat	Sys	Model	Interference
$K^{*0}\pi^+, K^{*0} \rightarrow K^+\pi^-$	10.3	± 1.2	± 0.7	$^{+0.4}_{-2.5}$	± 0.6
“higher K^{*0} ” $\pi^+, K^{*0} \rightarrow K^+\pi^-$	25.1	± 2.0	± 2.9	$^{+9.4}_{-0.5}$	± 4.9
$\bar{D}^0\pi^+, \bar{D}^0 \rightarrow K^+\pi^-$	184.6	± 3.2	± 9.7	-	-
$\rho^0 K^+, \rho^0 \rightarrow \pi^+\pi^-$	3.9 (< 6.2)	± 1.2	$^{+0.3}_{-0.6}$	$^{+0.3}_{-3.2}$	± 1.2
$f_0(980)K^+, f_0 \rightarrow \pi^+\pi^-$	9.2	± 1.2	± 0.6	$^{+1.2}_{-1.9}$	± 1.6
“higher f ” $K^+, f \rightarrow \pi^+\pi^-$	3.2 (< 12)	± 1.2	± 0.5	$^{+5.8}_{-2.4}$	± 1.5
Non resonant	5.2 (< 17)	± 1.9	$^{+0.8}_{-1.8}$	$^{+3.3}_{-7.5}$	± 6.4
$\chi_{c0}^0 K^+, \chi_{c0}^0 \rightarrow \pi^+\pi^-$	1.5	± 0.4	± 0.1	-	-

- Systematic errors are large since the exact nature of the contributions to the Dalitz Plot and their interferences are unknown
- “higher K^{*0} ” means any combination of $K_0^{*0}(1430)$, $K_2^{*0}(1430)$, $K_1^{*0}(1680)$
- “higher f ” means any combination of $f_2(1270)$, $f_0(1370)$, $f_0(1430)$

$B^\pm \rightarrow K^\pm \pi^\mp \pi^\pm$ Exclusive BF Conclusions

- Measured BFs with statistical significances $> 5\sigma$ for:
 - $B^\pm \rightarrow K^{*0}(896)\pi^\pm$
 - $B^\pm \rightarrow$ “higher K^{*0} ” π^\pm
 - $B^\pm \rightarrow f_0(980)K^\pm$
 - $B^\pm \rightarrow \chi_{c0}K^\pm$
 - $B^\pm \rightarrow D^0\pi^\pm$
- 90% CL upper limits for $B^\pm \rightarrow \rho^0 K^\pm$, $B^\pm \rightarrow$ “higher f ” K^\pm
- First tight limit on **non-resonant contribution**
- $K^{*0}(896)$ result larger than expected by **QCD factorization**
- Analysis documented in [hep-ex/0308065](https://arxiv.org/abs/hep-ex/0308065) and submitted to Physical Review D

Amplitude Analysis

- Greater statistics should allow the possibility of a **more thorough treatment**:
 - Full Dalitz Plot, or **Amplitude Analysis**
 - Complete treatment of **interferences** between the various resonances – should greatly reduce the large **systematic uncertainties** on the results for the sub decay modes
 - Measure **amplitude magnitudes and phases** → Branching Fractions, Charge Asymmetries and *CP* violation parameters
- This is what we are working on at the moment.
- Sian will tell you more in the next talk...