

The Relevance of Flavour Physics in the LHC Era

David Hitlin

Workshop on Flavour in the Era of the LHC

CERN

November 10, 2005

“The aim of physics is to find new terms in the Hamiltonian of the universe. Everything else is chemistry.”

M. Schwartz

By this stringent yardstick, most of us, while we strive to do physics, wind up doing chemistry

Nonetheless, flavour physics has historically been a rich source of physics, even by Mel's criterion

Does flavour physics in the LHC era meet this rigorous criterion?

What are the crucial experiments in the search for new terms in the flavour Hamiltonian (New Physics)?


Where is flavour physics now?

- The PEP-II/*BABAR* and KEKB/Belle *B* Factories, together with CLEO*c* and modern *K* decay experiments, have brought us to the precision measurement regime for many heavy quark and heavy lepton measurements
- CDF and DØ at Tevatron Run II are now producing physics and have unique capabilities in *B_s* mixing, rare decays and *b* baryon studies
- Bottom line: the CKM phase is consistent with being the source for all observed *CP*-violating phenomena
 - There must, however, be additional sources of *CP* violation
- Over the next few years, *CP* violation and rare decays in the flavour sector will be pursued with improved precision, and, equally important, with improved sophistication
 - *e.g.*, the best current methods for measurement of $\alpha(\phi_2)$ and $\gamma(\phi_1)$ were not those put forward in the initial phase

The Unitarity Triangle summarizes the salient features of Standard Model heavy flavour physics

$$V = \begin{pmatrix} V_{ud} & V_{us} & V_{ub} \\ V_{cd} & V_{cs} & V_{cb} \\ V_{td} & V_{ts} & V_{tb} \end{pmatrix} \cong \begin{pmatrix} 1 - \lambda^2/2 & \lambda & A\lambda^3(\bar{\rho} - i\bar{\eta}) \\ -\lambda & 1 - \lambda^2/2 & A\lambda^2 \\ A\lambda^3(1 - \bar{\rho} - i\bar{\eta}) & -A\lambda^2 & 1 \end{pmatrix}$$


Unitarity: $V_{ud}V_{ub}^* + V_{cd}V_{cb}^* + V_{td}V_{tb}^* = 0$




CP violation $\propto J = \Im[V_{ud}V_{cs}V_{us}^*V_{cd}^*] \simeq A^2\lambda^6\eta \sim 10^{-5}$, the Jarlskog invariant:

Let me count the ways

- There are many ways to overconstrain the Unitarity Triangle
- Useful constraints from B and K decays



Overconstraining the Unitarity Triangle



CP asymmetry from the interference between mixing and decay

$$A_{f_{CP}}(t) = \frac{\Gamma(\bar{B}^0(t) \rightarrow f_{CP}) - \Gamma(B^0(t) \rightarrow f_{CP})}{\Gamma(\bar{B}^0(t) \rightarrow f_{CP}) + \Gamma(B^0(t) \rightarrow f_{CP})}$$

$$A_{f_{CP}}(t) = S \cdot \sin(\Delta m \cdot t) - C \cdot \cos(\Delta m \cdot t)$$


$$S = \frac{2 \cdot \text{Im}(\lambda)}{1 + |\lambda|^2} \quad C = \frac{1 - |\lambda|^2}{1 + |\lambda|^2}$$

$$\lambda = \sqrt{\frac{M_{12}^* - \frac{i}{2}\Gamma_{12}^*}{M_{12} - \frac{i}{2}\Gamma_{12}}} \cdot \frac{\langle f_{CP} | H | \bar{B}^0 \rangle}{\langle f_{CP} | H | B^0 \rangle} = \frac{q}{p} \cdot \frac{\bar{A}_f}{A_f}$$

For a single decay amplitude: $|q/p|=1$


$$|\lambda|=1 \Rightarrow S = \text{Im}(\lambda), \quad C = 0$$

$$A_{f_{CP}}(t) = \text{Im}(\lambda) \cdot \sin(\Delta m \cdot t)$$




If there is New Physics
it could be in the decay
amplitude or the mixing
amplitude

$\sin 2\beta$ is now a precision measurement




$J/\psi, \psi(2S), \chi_{c1}, \eta_c$

$\bar{K}^0 \rightarrow K_S^0, K_L^0$
 $\bar{K}^{*0} \rightarrow K_S^0 \pi^0$




$$\sin 2\beta = +0.722 \pm 0.040 \pm 0.023$$



$$\sin 2\phi_1 = +0.652 \pm 0.039 \pm 0.020$$


Unitary Triangle 2005




$|V_{td}/V_{ts}|$ from $B \rightarrow \rho\gamma$ vs. $B \rightarrow K^*\gamma$

Is there New Physics in $B^0\bar{B}^0$ mixing?

- Assume: New Physics in tree decays - negligible
- Define $r_d^2 \exp(2i\theta_d) \equiv 1 + h e^{i\sigma} \equiv \langle B^0 | \mathcal{H}^{\text{full}} | \bar{B}^0 \rangle / \langle B^0 | \mathcal{H}^{\text{SM}} | \bar{B}^0 \rangle$
- Use $|V_{ub}/V_{cb}|$, \mathcal{A}_{DK} , $S_{\psi K}$, $S_{\rho\rho}$, Δm_{B_d} , \mathcal{A}_{SL}
- Fit to η , ρ , r_d , θ_d (or h , σ)
- Find whether $h \neq 0$ ($r_d \neq 1$) is allowed



Nir




Agashe, et al.

For arbitrary phase, $h = |A_{\text{NP}}/A_{\text{SM}}| \sim 0.2 \pm 0.2$



B_s mixing limits

□ New CDF/DØ results




Limit 14.5 \rightarrow 16.6 ps⁻¹
sensitivity 18.3 \rightarrow 20.0 ps⁻¹

LHCb sensitivity to $B_s^0 \bar{B}_s^0$ mixing




In 1 year, $\geq 5\sigma$ observation of B_s^0 oscillations up to $\Delta M_s = 68 \text{ ps}^{-1}$

Is there New Physics in $B_s^0 \bar{B}_s^0$ mixing?



Current



If $\Delta m_s = (18.3 \pm 0.3) \text{ps}^{-1}$


Agashe, Papucci, Perez, Pirjol

ϕ_s

Mixing-induced CP violation: phase mismatch $\phi_s - 2\phi_D \approx \phi_s \neq 0, \pi$

“first mix, then decay”

• $\phi_s^{\text{SM}} \equiv 2 \arg [V_{ts}^* V_{tb}] \approx -2\beta_s = \mathcal{O}(-0.04)$ rad



→ CP-asymmetry directly measures $\phi_s = \mathcal{O}(-0.04)$ rad (for given $\eta_{f_{\text{CP}}}$)

$$\mathcal{A}_{\text{CP}}(t) = \frac{-\eta_{f_{\text{CP}}} \sin(\phi_s) \sin(\Delta M_s t)}{\cosh\left(\frac{\Delta\Gamma_s t}{2}\right) - \eta_{f_{\text{CP}}} \cos(\phi_s) \sinh\left(\frac{\Delta\Gamma_s t}{2}\right)}$$

Channels	$\sigma(\phi_s)$ [rad]	Weight $(\sigma/\sigma_i)^2$ [%]
$B_s^0 \rightarrow J/\psi \eta(\gamma \gamma)$	0.112	6.4
$B_s^0 \rightarrow J/\psi \eta(\pi^+ \pi^- \pi^0)$	0.148	3.6
$B_s^0 \rightarrow \eta_c \phi$	0.106	7.1
Combined three pure CP eigenstates channels	0.068	17.1
$B_s^0 \rightarrow J/\psi \phi$	0.031	82.9
Combined all four CP eigenstates channels	0.028	100.0

With 10 fb^{-1} (5 years): $\sigma(\phi_s) \sim 0.013$ rad → $\sim 3\sigma$ for $\phi_s = -0.04$ rad (SM)

Constraints on New Physics


- Even at current precision, the *grosso modo* agreement of flavour physics experimental results with Standard Model predictions places very meaningful limits on physics beyond the Standard Model
- In the presence of New Physics, many of these simple relations can be modified
- This has spawned a minor phenomenological industry that has provided guidance for the current experiments and a roadmap for future investigations
 - Where to look for New Physics?
 - *FCNC processes*
 - $b \rightarrow s$ vertices: $\Delta F=1$, $\Delta F=2$
 - CP violation, mixing, rare b and τ decays
 - Precision measurements in K , D and B decays

The LHC era begins

- What will remain to be done in flavour physics when the current programs at *BABAR*, Belle, CDF and DØ are over?
 - Overconstrained tests of the CKM matrix to the level of precision warranted by theoretical uncertainties
 - Searches, and, quite likely, measurements of deviations from the Standard Model that will be crucial to an understanding of New Physics uncovered at the LHC
- What facilities will we have?
 - LHC: ATLAS, CMS, LHC*b*
 - Linear collider (ILC/CLIC)?
 - A Super *B* Factory?
 - Rare kaon experiments?
 - τ /charm collider
 - Fixed target charm experiment?
- Which facilities beyond the LHC are most crucial to flavour studies?

An Endangered Species Act for heavy quarks in the US?

2003



2005





2005



2005



2006-2008



Data samples (b) at the start* of the LHC era

- *BABAR* and Belle will each have collected a total data sample of approximately 1 ab^{-1} by ~ 2008
 - $2 \text{ ab}^{-1} = 2 \times 10^9$ produced $B\bar{B}$ pairs
- The Tevatron Run II experiments CDF and DØ will each have collected $4\text{-}8 \text{ fb}^{-1}$ by ~ 2009
 - Premier objective is a measurement of $B_s^0 \bar{B}_s^0$ mixing

How can flavour physics complement the LHC program?

- The Higgs will likely come first
- The search for evidence of physics beyond the Standard Model will then take center stage
 - There have been many exercises showing how, with sufficiently large data samples and concomitant improved precision, heavy quark and τ measurements can uniquely discriminate between SUSY breaking models, extra dimension scenarios, *etc.*
 - These need sharpening, on both sides: high p_t and flavour physics
- It is possible that the LHC will find the Standard Model Higgs and nothing else
 - In this situation, precision flavour physics may be the best way we have to search for physics beyond the Standard Model


Strategies

- Improve measurements of CKM unitarity triangle measurements to the precision warranted by theory
 - CP asymmetries in tree-dominated processes
 - Differences in CP asymmetries between tree and penguin-dominated processes

- Search for/measure FCNC processes, highly suppressed in the Standard Model, in which New Physics may show up
 - Generation 3 to generation 2 ($b \rightarrow s$, $\tau \rightarrow \mu$) transitions are especially promising
 - B_s oscillations
 - $b \rightarrow s \gamma$ branching fraction, A_{CP}
 - $A_{FB}(b \rightarrow s \ell^+ \ell^-)$
 - $B_s \rightarrow \mu^+ \mu^-$

Precision of α , $\sin 2\beta$, γ in this decade

- Through ~ 2008 (and beyond), precision of UT angle measurements will continue to be statistics limited



- α , γ methods shown run into model problems at $2 ab^{-1}$ level
 - With higher statistics, other methods may be better
 - e.g.* for α , use $B \rightarrow \pi\pi$ + isospin analysis instead of $B \rightarrow \rho\rho$


LHCb (1 year @ 2×10^{32}) (Schneider)

$$\sigma(\alpha) = 10^0$$

$$\sigma(\gamma) = 5^0 (B^\pm \rightarrow D^0 K^\pm)$$

$$\sigma(\sin 2\beta) = 0.02$$

Global CKM fit: 2008



$\sigma(V_{ub}) = 6.5\%$

$\sigma(\Delta m_s) = 5\%$

$\sigma(\sin 2\beta) = 0.019 \quad \sigma(\alpha) = 8^\circ \quad \sigma(\gamma) = 10^\circ$

What is the ultimate possible UT precision?


- Theoretical limits (continuum methods)
 - Many measurements will not be theory-limited for quite some time

Ligeti:


Measurement (in SM)	Theoretical limit	Present error
$B \rightarrow \psi K_S$ (β)	$\sim 0.2^\circ$	1.6°
$B \rightarrow \phi K_S, \eta^{(\prime)} K_S, \dots$ (β)	$\sim 2^\circ$	$\sim 10^\circ$
$B \rightarrow \pi\pi, \rho\rho, \rho\pi$ (α)	$\sim 1^\circ$	$\sim 15^\circ$
$B \rightarrow DK$ (γ)	$\ll 1^\circ$	$\sim 25^\circ$
$B_s \rightarrow \psi\phi$ (β_s)	$\sim 0.2^\circ$	—
$B_s \rightarrow D_s K$ ($\gamma - 2\beta_s$)	$\ll 1^\circ$	—
$ V_{cb} $	$\sim 1\%$	$\sim 3\%$
$ V_{ub} $	$\sim 5\%$	$\sim 15\%$
$B \rightarrow X\ell^+\ell^-$	$\sim 5\%$	$\sim 20\%$
$B \rightarrow K^{(*)}\nu\bar{\nu}$	$\sim 5\%$	—
$K^+ \rightarrow \pi^+\nu\bar{\nu}$	$\sim 5\%$	$\sim 70\%$
$K_L \rightarrow \pi^0\nu\bar{\nu}$	$< 1\%$	—

Ultimate UT precision - II

- Unitarity triangle using unquenched lattice results



Current precision




"Ultimate" precision

Okamoto

Ultimate UT precision - III


- Prediction for SuperKEKB at 50 ab^{-1} (~ 2020)
- LHCb measurements will also be pertinent (or, they may be the only measurements)



- Rare K decay constraints

	short-distance (e.w.) contrib. to the total rate $(\Gamma - \Gamma_{\text{no s.d.}}) / \Gamma$	present irreducible th. error on the s.d. amplitude extracted from BR only	total BR within SM (central value)
$K_L \rightarrow \pi^0 \nu \nu$	> 99%	1%	8×10^{-11}
$K^+ \rightarrow \pi^+ \nu \nu$	88%	3%	3×10^{-11}
$K_L \rightarrow \pi^0 e^+ e^-$	38%	15% → Dalitz $\sim 10\%$	3.5×10^{-11}
$K_L \rightarrow \pi^0 \mu^+ \mu^-$	28%	30% → Dalitz $\sim 15\%$	1.5×10^{-11}

- The **real** motivation for precision measurements of rare K decay branching fractions is the sensitivity to New Physics




- The same is true for high precision B and τ decay studies

After the LHC finds New Physics

- LHC, by the mass difference and threshold methods, will measure SUSY masses to a precision of 5 to 10 GeV
- ILC or CLIC, **a decade later**, will dramatically improve errors, by, e.g., measuring slepton masses with a threshold scan to a precision ~ 100 MeV with 10 fb^{-1} per point


	LHC	LHC+ILC
$\Delta m_{\tilde{\chi}_1^0}$	4.8	0.19
$\Delta m_{\tilde{l}_R}$	4.8	0.34
$\Delta m_{\tilde{\chi}_2^0}$	4.7	0.24
$\Delta m_{\tilde{q}_L}$	8.7	4.9
$\Delta m_{\tilde{b}_1}$	13.2	10.5




Correlations allow model discrimination (without ILC)

- In the meantime it is important to clarify the flavour structure of New Physics discoveries with high statistics K , B and τ experiments
- These studies can uniquely discriminate between SUSY models

Many SM extensions yield measurable effects in B physics



Heavy flavour studies complement LHC direct searches



Experimental signatures of extended flavour structure

- FCNC processes, LFV processes and CPV in loop decays
- There are many clean measurements that are diagnostic of models
 - New CP violating phase in $b \rightarrow sqq$: $CPV(t)$ in $B^0 \rightarrow \phi K_{s'}, \eta' K_{s'}, K_s K_s K_s$
 - Right-handed current in $b \rightarrow s\gamma$: $CPV(t)$ in $B^0 \rightarrow K_s \pi^0 \gamma$
 - Lepton flavour violation in τ decays : $\mathcal{B}(\tau \rightarrow \mu\gamma)$
 - Charged Higgs in tree diagram : $\mathcal{B}(B \rightarrow D\tau\nu)/\mathcal{B}(B \rightarrow D\mu\nu)$
 - $b \rightarrow d\gamma, A_{FB}$ in $b \rightarrow sl^+\ell^-, B \rightarrow K^*\nu\nu, B \rightarrow \ell\nu, \dots\dots\dots$
 - B_s mixing


- Some measurements are unique to hadron experiments, some to e^+e^- , and some can be done in both environments

Many rare processes are sensitive to New Physics

Mode	Standard Model Branching Fraction \mathcal{B}	New Physics Effects
$b \rightarrow s\gamma$	$\sim 3 \times 10^{-4}$	$\mathcal{B}, A_{CP}, A_{FB}$
$b \rightarrow sg$	$\sim 10^{-5}$ per mode	A_{CP}
$B \rightarrow X\ell\ell$	$\sim 10^{-6}$ per mode	$\mathcal{B}, A_{CP}, A_{FB}$
$B \rightarrow X\nu\nu$	$\sim 10^{-6}$ per mode	\mathcal{B} up to 10^{-5} /mode
$D \rightarrow X\ell\ell$	$\sim 10^{-6}$ per mode	\mathcal{B} up to 10^{-5} /mode
$B \rightarrow \tau\nu$	$\sim 10^{-4}$	Close to SM sensitivity
$\tau \rightarrow \ell\gamma$	$\sim 10^{-40}$	\mathcal{B} up to 10^{-8}
$B \rightarrow \ell\ell$	$< 10^{-11}$	\mathcal{B} up to 10^{-5}
$D \rightarrow \ell\ell$	$< 10^{-9}$	\mathcal{B} up to 10^{-6}

The observed pattern of effects is diagnostic of the mechanism of SUSY breaking or the type of extra dimension model

Flavour physics provides a roadmap




What can a Super B Factory bring to the party?

- Flavour physics, whether b , c or τ decays, provides
 - Sensitivity to New Physics (SUSY, extra dimensions,)
 - New CP phases, non-Standard Wilson coefficients,
 - Strong and unique constraints on models of SUSY breaking
 - $\mathcal{B}(b \rightarrow s \gamma)$ has already ruled out a host of New Physics models
 - Super B specific measurements
 - CPV in $b \rightarrow s$
 - FCNC ($Kll, K\nu\nu$
 - LFV (τ decays)
 - Higgs mediation ($B \rightarrow \tau\nu, B \rightarrow D\tau\nu$, etc.)
 - Precision CKM Unitarity Triangle parameters
 - Measure $\alpha, \beta, \gamma, V_{ub}, V_{cb}, \Delta m_d$ to the limit of theoretical precision (see above)


- There is, of course, some overlap with LHC **b**

Probes of new physics


- In the Standard Model we expect the same value for “ $\sin 2\beta$ ” in $b \rightarrow c\bar{c}s$, $b \rightarrow c\bar{c}d$, $b \rightarrow s\bar{s}s$, $b \rightarrow d\bar{d}s$ modes, but different SUSY models can produce **different asymmetries**
- Since the penguin modes have branching fractions one or two orders of magnitude less than tree modes, great deal of luminosity is required to make these measurements to meaningful precision



$$\lambda_{tree} = \frac{q}{p} \frac{\bar{A}}{A} = \eta \frac{V_{tb}^* V_{td} V_{cb} V_{cs}^*}{V_{tb} V_{td}^* V_{cb}^* V_{cs}} = (-1) e^{-2i\beta}$$









$$\lambda_{penguin} = \frac{q}{p} \frac{\bar{A}}{A} = \eta \frac{V_{tb}^* V_{td} V_{tb} V_{ts}^*}{V_{tb} V_{td}^* V_{tb}^* V_{ts}} = (-1) e^{-2i\beta}$$



$$\lambda = e^{i(2\beta + \phi^{SUSY})} \frac{|\bar{A}|}{|A|} \Rightarrow S_{\phi K} = \sin(2\beta + \phi^{SUSY})$$

In general $S_{J/\psi K_S} \neq S_{\phi K_S}$; $C_{J/\psi K_S} \neq C_{\phi K_S}$

There are penguin and tree corrections to $b \rightarrow ss\bar{s}, sdd\bar{d}$ modes


$b \rightarrow ss\bar{s}, sdd\bar{d}$ mode	\mathcal{B} $\times 10^{-6}$		
$B^0 \rightarrow \phi K_S^0$	4	$\propto V_{tb}V_{ts}^* \sim \lambda^2$ 	$\propto V_{ub}V_{us}^* \sim \lambda^4 R_u e^{-i\gamma}$ 
$B^0 \rightarrow \eta' K_S^0$	29	$\propto V_{tb}V_{ts}^* \sim \lambda^2$ 	$\propto V_{ub}V_{us}^* \sim \lambda^4 R_u e^{-i\gamma}$ 
$B^0 \rightarrow \pi^0 K_S^0$	20	$\propto V_{tb}V_{ts}^* \sim \lambda^2$ 	$\propto V_{ub}V_{us}^* \sim \lambda^4 R_u e^{-i\gamma}$ 

$B^0 \rightarrow J/\psi K_S^0$ 440

Issues: triggering, charged/neutral vertexing,

There is as yet no evidence for New Physics

from CP asymmetry measurements in $b \rightarrow s$ transitions




Kirkby and Nir, PDG

As we press the search, we must understand

- The required and achievable experimental precision
- The reliability of theoretical calculations

How good are the SM predictions?



Two body:
Beneke, PLB 620 (2005) 143

blue = ± 1 sigma
grey = full range

Calculations within
framework of QCD
factorization

Three body:
Cheng, Chua & Soni,
hep-ph/0506268

$$\Delta = S_{penguin} - \sin 2\beta$$

squark mass matrix (d sector)

Flavour structure:
SuperB, LHCb


$$\begin{pmatrix}
 m_{\tilde{d}_L}^2 & m_d(A_d - \mu \tan \beta) & (\Delta_{12}^d)_{LL} & (\Delta_{12}^d)_{LR} & (\Delta_{13}^d)_{LL} & (\Delta_{13}^d)_{LR} \\
 & m_{\tilde{d}_R}^2 & (\Delta_{12}^d)_{RL} & (\Delta_{12}^d)_{RR} & (\Delta_{13}^d)_{RL} & (\Delta_{13}^d)_{RR} \\
 & & m_{\tilde{s}_L}^2 & m_s(A_s - \mu \tan \beta) & (\Delta_{23}^d)_{LL} & (\Delta_{23}^d)_{LR} \\
 & & & m_{\tilde{s}_R}^2 & (\Delta_{23}^d)_{RL} & (\Delta_{23}^d)_{RR} \\
 & & & & m_{\tilde{b}_L}^2 & m_b(A_b - \mu \tan \beta) \\
 & & & & & m_{\tilde{b}_R}^2
 \end{pmatrix}$$

Mass spectrum: LHC, LC

Assuming all Δ 's small and squarks nearly degenerate, we can use mass insertion approximation (MIA):

$$(\delta_{ij}^d)_{AB} = \frac{(\Delta_{ij}^d)_{AB}}{\tilde{m}^2}$$

Flavour mixing parametrization
 $b \rightarrow s$ (23), $b \rightarrow d$ (13)




Off diagonal terms can provide unique information on CP phases

The scale of New Physics


- Mass insertion approximation: model-independent ($\delta_{LL,RR}$)

ϕ_{13} mass insertion



$\Delta A_{CP}(J/\psi K_S - \pi^0 K_S)$

ϕ_{23} mass insertion




$\Delta A_{CP}(J/\psi K_S - \phi K_S)$







after Ciuchini, Franco, Martinelli, Masiero, & Silvestrini

How much integrated luminosity ?

- 50 ab^{-1} is required for CP asymmetry measurements in rare penguin modes and for rare branching fractions





New Physics effects in the **MMSM/THDM**



Time dependent CP asymmetry in $B \rightarrow X_s \gamma$

- Theoretically clean, but A_{CP} is very small in the Standard Model, since the photon is polarized, and the final state is almost flavor specific
 \Rightarrow Helicity suppression:
 $\sim m_s/m_b$

- Requires vertex reconstruction with K_S alone



SuperKEKB study:

$$\sigma(A_{CP}^{mix}(B \rightarrow K^* \gamma, K^* \rightarrow K_S \pi^0)) = 0.04$$

$$\sigma(A_{CP}^{dir}(B \rightarrow X_s \gamma)) = 0.005$$

for 50 ab^{-1}

Asymmetric B factories provide a uniquely clean environment

- High (30%) and well-measured flavour-tagging efficiency

- B decays with neutrinos

$$B \rightarrow D\tau\nu, \tau\nu, ul\nu, \dots$$

Charged Higgs

V_{ub}

- B decays with γ, π^0



$$B \rightarrow X_s\gamma, \pi^0\pi^0, \dots$$

- B vertex reconstruction with K_S^0 alone

$$B \rightarrow K_S^0\pi^0, \rightarrow K_S^0\pi^0\gamma, \dots$$


direct CPV

$\alpha(\phi_2)$ isospin analysis




A_{CP}^{mix} in four SUSY models

Direct CPV



$m_{\tilde{g}} (GeV)$

Mixing CPV





$m_{\tilde{g}} (GeV)$

50ab⁻¹

Goto, Okada, Shimizu, Shindou, Tanaka



$$b \rightarrow s l^+ l^-, B \rightarrow K l^+ l^-, B \rightarrow K^* l^+ l^-$$

- Uniquely sensitive tests of Wilson coefficients



$$b \rightarrow s \ell^+ \ell^-, B \rightarrow K \ell^+ \ell^-, B \rightarrow K^* \ell^+ \ell^-$$

- Exclusive decays affected by larger uncertainties
- Inclusive decays are better: only in e^+e^-
- Use of ratios: A_{FB} , ACP , $e^+e^-/\mu^+\mu^-$




$$b \rightarrow s \ell^+ \ell^-, B \rightarrow K \ell^+ \ell^-, B \rightarrow K^* \ell^+ \ell^-$$

Precision




Sup




LHC

$-A_{FB}$ vs q^2

SuperB factory with 50



C_{10}^{NP}




A_{FB} K*ll with 5/ab

- 2 fb⁻¹ : (4.0 ± 1.2) GeV² with
 - 10 fb⁻¹ : (4.0 ± 0.5) GeV²
- ⇒ 13% error on C_7^{Eff}/C_9^{Eff}

$$b \rightarrow s \ell^+ \ell^-, B \rightarrow K \ell^+ \ell^-, B \rightarrow K^* \ell^+ \ell^-$$

- Extra dimension models
 - Randall-Sundrum example (Rizzo)
 - Zero crossing moves with mass of lightest KK graviton







(More) model-independent analysis of NP couplings

New Physics scalar and pseudoscalar couplings can result in a difference in

$$R_{K,K^*,x_S} = \frac{\mathcal{B}(B \rightarrow (K, K^*, x_S) e^+ e^-)}{\mathcal{B}(B \rightarrow (K, K^*, x_S) \mu^+ \mu^-)}$$

from unity

R_{K,K^*,x_S} is correlated with the (unobserved) rate $\mathcal{B}(B_s \rightarrow \mu^+ \mu^-)$



Hiller and Krüger


(More) model-independent analysis of NP couplings

New Physics scalar and pseudoscalar couplings can result in a difference in


$$R_{K,K^*,x_s} = \frac{\mathcal{B}(B \rightarrow (K, K^*, x_s) e^+ e^-)}{\mathcal{B}(B \rightarrow (K, K^*, x_s) \mu^+ \mu^-)}$$

from unity

R_{K,K^*,x_s} is correlated with the (unobserved) rate $\mathcal{B}(B_s \rightarrow \mu^+ \mu^-)$




Hiller and Krüger




Koppenburg

Prospects

- R_K can be measured very well at a Super B factory
- It would appear to be difficult at LHCb due to difficulty with triggering on electrons, but a new study indicates that if the trigger can be modified, there is good sensitivity
- A 10% measurement of R_K is possible with 2.5 ab^{-1} at a Super B factory or with 2 fb^{-1} at LHCb with an electron trigger



Rare B decays: $B_s^0 \rightarrow \mu^+ \mu^-$




□ Standard Model $\mathcal{B} \sim 3.5 \times 10^{-9}$

■ Current limit (CDF+DØ) $< 1.5 \times 10^{-7}$ @ 95% CL

□ Should be observed at LHC

	1 year	$B_s \rightarrow \mu^+ \mu^-$ signal (SM)	$b \rightarrow \mu, b \rightarrow \mu$ background	Inclusive bb background
LHCb	2 fb^{-1}	17	< 100	< 7500
ATLAS	10 fb^{-1}	7	< 20	
CMS (1999)	10 fb^{-1}	7	< 1	



Excluded!

$B \rightarrow \tau \nu$




Standard Model prediction:

$$BF(B^+ \rightarrow \tau^+ \nu_\tau) = 1.2 \times 10^{-4} \left(\frac{f_B}{200 \text{ MeV}} \right)^2 \left(\frac{V_{ub}}{0.004} \right)^2$$

can be modified by an H^+ at large $\tan \beta$

$$BF(B^+ \rightarrow \tau^+ \nu_\tau) = 1.3^{+1.0}_{-0.9} \times 10^{-4}$$

($< 2.6 \times 10^{-4}$ at 90% C.L.)





$B \rightarrow \tau \nu$ limit on H^+ mass

$$\mathcal{B}(B \rightarrow \tau \nu) = \mathcal{B}(B \rightarrow \tau \nu)_{\text{SM}} \times r_H,$$

$$r_H = \left(1 - \frac{m_B^2}{m_H^2} \tan^2 \beta \right)^2$$


95% CL excluded region
at 5 ab^{-1} (if $\mathcal{B}_{\text{obs}} = \mathcal{B}_{\text{SM}}$)




Iijima

$$B^+ \rightarrow \bar{D}^0 \tau \nu_\tau$$

- Standard Model: $\mathcal{B} = 8 \times 10^{-3}$




- For 50 ab^{-1} : $\sigma \mathcal{B} / \mathcal{B} = 2.5\%$



SuperKEKB


$\tau \rightarrow l\gamma$ and mSUGRA



The pattern of deviation from the SM values is diagnostic

Model	B_d Unitarity	Time-dep. CPV	Rare B decay	Other signals
mSUGRA (moderate $\tan \beta$)	-	-	-	-
mSUGRA (large $\tan \beta$)	B_d mixing	-	$B \rightarrow (D)^* \tau \nu$ $b \rightarrow s \ell^+ \ell^-$	$B_s \rightarrow \mu \mu$ B_s mixing
SUSY GUT with ν_R	-	$B \rightarrow \phi K_S$ $B \rightarrow K^* \gamma$	-	B_s mixing τ LFV, n EDM
Effective SUSY	B_d mixing	$B \rightarrow \phi K_S$	$A_{CP}(b \rightarrow s \gamma)$ $b \rightarrow s \ell^+ \ell^-$	B_s mixing
KK graviton exchange	-	-	$b \rightarrow s \ell^+ \ell^-$	-
Split fermions in large extra dimensions	B_d mixing	-	$b \rightarrow s \ell^+ \ell^-$	$K^0 \bar{K}^0$ mixing $D^0 \bar{D}^0$ mixing
Bulk fermions in warped extra dimensions	B_d mixing	$B \rightarrow \phi K_S$	$b \rightarrow s \ell^+ \ell^-$	$D^0 \bar{D}^0$ B_s mixing mixing
Universal extra dimensions	-	-	$b \rightarrow s \ell^+ \ell^-$ $b \rightarrow s \gamma$	$K \rightarrow \pi \nu \nu$

Flavour physics is a DNA chip for New Physics



The role of hadronic experiments

- Measure $\mathcal{B}(K^+ \rightarrow \pi^+ \nu \bar{\nu})$ and $\mathcal{B}(K_L^0 \rightarrow \pi^0 \nu \bar{\nu})$
- Search for LFV in μ to e conversion, $\mu \rightarrow e \gamma$

- Measure B_s mixing
- Measure CPV in $J/\psi \phi$
- Measure $\mathcal{B}(B_s \rightarrow \mu \mu)$ at SM level or above
- Measure A_{FB} in exclusive $b \rightarrow s \ell \ell$
- Measure $\mathcal{B}(B \rightarrow K e e) / \mathcal{B}(B \rightarrow K \mu \mu)$?
- Improve measurements of α, β, γ

The role of a Super B Factory (if one exists)


- Measure $\alpha, \beta, \gamma, V_{ub}, V_{cb}, \Delta m_d$ to the limit of theoretical precision
- Measure CPV in $b \rightarrow sss$ decays to sufficient precision to ascertain a pattern
- Measure A_{FB} in inclusive and exclusive $b \rightarrow sl\ell$ decays
- Measure $\mathcal{B}(B \rightarrow Kee)/\mathcal{B}(B \rightarrow K\mu\mu)$
- Measure $B \rightarrow \tau\nu$
- Search for right handed couplings in $b \rightarrow s\gamma$
- Search for NP effects in $B \rightarrow VV$ decay
- Search for CPV in $b \rightarrow s\gamma$ and $b \rightarrow sl\ell$ decays
- Search for LFV: $\tau \rightarrow \mu\gamma$ decay
- Search for charged Higgs in $B \rightarrow D\tau\nu$ decay

Will there be a Super B Factory?

- Potential upgrades of PEP-II, KEKB
 - SuperPEP-II design aimed at $\mathcal{L}=7 \times 10^{35}$
 - SLAC is now out of the local HEP accelerator business
 - SuperKEKB design has $\mathcal{L}=4 \times 10^{35}$ with 50 ab^{-1} by 2020
 - Proposed to ministry; awaits a decision
- Another approach is being considered
 - Raimondi has suggested revisiting the recirculating linear collider scheme proposed in the '80's by Amaldi and Coignet, in the light of modern accelerator physics techniques
 - The luminosity goal is $\mathcal{L} \sim 10^{36}$
 - The idea appears quite promising
 - Linear collider-related R&D over the past two decades has addressed many of the technical issues
 - INFN is actively considering an LCSuper B based on an international collaborative effort
 - A workshop will be held on Nov 11/12 at Frascati

<http://www.lnf.infn.it/conference/superbf05/>

One and one half Super B Factory concepts



Conclusions

- Heavy flavour physics, in all its guises, can play a significant role in deepening our understanding of the Standard Model, and, should New Physics be found at LHC, it provides unique tools for probing the flavour structure of the new particles (squarks?)
- If the scale of New Physics is < 1 TeV, as motivated by our current understanding of the Higgs mechanism, there will be measurable effects in the heavy quark/heavy lepton sector
 - LHC will yield information on masses and couplings of new particles by direct production
 - Information on squark off-diagonal couplings requires detailed studies of heavy quark decays at Super B Factories or dedicated hadronic experiments
- The effects of new physics loops can be seen in rare decay branching fractions and kinematic distributions and in CP -violating asymmetries in channels with very small branching fractions

Conclusions - II

- The new generation of experiments at hadronic accelerators will doubtless extend the fruitful programs of the current *B* Factories
- It is important that other approaches be followed as well:
 - A Super *B* Factory can, in the next decade, provide **unique high precision measurements** as well as results complementary to those of hadronic experiments
 - Rare K decay experiments
 - Searches for lepton flavor violation
- Flavour physics has a glorious history and remains relevant as a tool for understanding New Physics at the LHC
- This timely series of workshops can sharpen our understanding of what is needed at LHC and at other facilities to deeply probe the structure of New Physics, in particular the flavour structure

