



***Status and Prospects of
Rare K^\pm and K_L Decays
from NA48***

Rainer Wanke

Institut für Physik, Universität Mainz

Kaon Mini Workshop

CERN, May 2, 2005

■ K^\pm Decays:


- $K^\pm \rightarrow \pi^\pm \pi^\mp e^\pm \nu(\bar{\nu})$ (K_{e4}^+) branching ratio and form factors
- $K^\pm \rightarrow \pi^\pm e^+ e^-$ and $K^\pm \rightarrow \pi^\pm \mu^+ \mu^-$ decays
- Branching ratio and form factors of $K^\pm \rightarrow \pi^\pm \pi^0 \gamma$
- Branching ratio and form factors of $K^\pm \rightarrow \pi^\pm \gamma \gamma$

■ $K_{L,S}$ Decays:

- Precise measurements of $K_L \rightarrow \pi^+ \pi^- / \pi^0 \pi^0$ branching ratios
- Measurement of $K_S \rightarrow \pi^+ \pi^- \pi^0$.
- Search for $K_S \rightarrow 3\pi^0$.

Most analyses: Not yet finished

⇒ **Event numbers and statistical reach shown, but few results.**




Rare K^\pm Decays

NA48/2 K^\pm Data Taking

NA48/2 experiment in 2003/2004:

- **2003:** ~ 50 days of data taking: Super samples 0, 1 – 3
 \Rightarrow Data shown in this talk (partly without SS 0).
- **2004:** ~ 60 days of data taking: Super samples 4 – 8
 \Rightarrow Data *not* shown here, but simple to add.



Trigger:

- All 3-track events ($\sim 98\%$ efficient)
- 1-track events with μ -veto and $(p_K - p_\pi)^2 \gg m_{\pi^0}^2$ to reject $K^\pm \rightarrow \pi^\pm \pi^0$.

Measurement of $K^+ \rightarrow \pi^+ \pi^- e^+ \nu$ (K_{e4}^+)

- Physics interest: (Same as cusp-effect in $K^\pm \rightarrow \pi^\pm \pi^0 \pi^0$)
 - Low energy $\pi\pi$ scattering unambiguously predicted from Chiral Perturbation Theory first principles.
 - s -wave isospin zero $\pi\pi$ scattering length a_0^0 can be determined from form factors in K_{e4}^+ decays.
 - ⇒ Predicted to $a_0^0 = 0.220 \pm 0.005$ in ChPT.
(Colangelo, Gasser, Leutwyler, 2001)
- Previous measurements:
 - Geneva-Saclay (1977): 30 000 events
 - Brookhaven E865 (2001): 400 000 events
 - ⇒ $a_0^0 = 0.216 \pm 0.013$.
 - NA48 (2005, prel.): $|a_0^0 - a_2^0|$ from $K^\pm \rightarrow \pi^\pm \pi^0 \pi^0$
 - ⇒ $|a_0^0 - a_2^0| = 0.281 \pm 0.021$.

Measurement of $K^+ \rightarrow \pi^+ \pi^- e^+ \nu$ (K_{e4}^+)

- Events fully reconstructed by using known kaon momenta.

(Well, $p = 60 \text{ GeV}/c$ along z -axis.)

⇒ About 340 000 events in SS 1–3.

- **Background** mainly from

$K^+ \rightarrow \pi^+ \pi^+ \pi^-$ and

$K^+ \rightarrow \pi^+ \pi_{\text{Dalitz}}^0$ with


$\pi \leftrightarrow e$ mis-identification.

⇒ Use of neural network for e/π separation.

- Total background: 0.4%







Determined from wrong-sign

$\pi^+ \pi^+ e^-$ data events.



Measurement of $K^+ \rightarrow \pi^+ \pi^- e^+ \nu$ (K_{e4}^+)

Cabibbo-Maksymowicz variables: (Background in red)



Measurement of $K^+ \rightarrow \pi^+ \pi^- e^+ \nu$ (K_{e4}^+)

Status and Prospects for K_{e4}^+ :

- About **340 000 events** in SS 1–3 2003.
 - ⇒ Expected statistical error on $a_0^0 \sim \pm 0.01$.
 - ⇒ Similar precision as cusp analysis in $K^+ \rightarrow \pi^+ \pi^0 \pi^0$, but complementary & no theoretical uncertainties.
- Backgrounds are tiny, systematics are under study.
- Analysis underway, but a lot of work.

Further K_{l4}^+ Analyses:

- About **15 000 events** of $K^\pm \rightarrow \pi^0 \pi^0 e^\pm \nu(\bar{\nu})$ in SS 1–3.
- Also $K^\pm \rightarrow \pi^+ \pi^- \mu^\pm \nu(\bar{\nu})$ under investigation.

Both measurements have to fight large backgrounds.

Measurement of

$$K^\pm \rightarrow \pi^\pm e^+ e^- \text{ and } K^\pm \rightarrow \pi^\pm \mu^+ \mu^-$$

■ Physics interest:


- FCNC, suppressed by GIM mechanism.
- Amplitude dominated by long-distance contributions (one-photon “bremsstrahlung” diagrams).
⇒ Can be extracted from form factor measurements!
- Potentially interesting channels for CP violation between $\Gamma(K_{\pi ll}^+)$ and $\Gamma(K_{\pi ll}^-)$.

■ Previous measurements:

- $K^\pm \rightarrow \pi^\pm e^+ e^-$: BNL E865 found 10500 events
⇒ $\text{Br} = (2.94 \pm 0.05 \pm 0.14) \times 10^{-7}$
(Also form factor measurement.)
- $K^\pm \rightarrow \pi^\pm \mu^+ \mu^-$: Several experiments, in total 800 events.
⇒ $\text{Br} = (0.81 \pm 0.14) \times 10^{-7}$

Measurement of $K^\pm \rightarrow \pi^\pm e^+ e^-$



Data: SS0-3



- More than **4000 events** for $m_{ee} > 140 \text{ MeV}/c^2$ in 2003 data.
- Background very small ($< 1\%$).

Measurement of $K^\pm \rightarrow \pi^\pm e^+ e^-$


■ Form Factor:



■ Status and Prospects:

- **Statistical error** on branching fraction $\Delta\text{Br} \leq \pm 0.05 \times 10^{-7}$.
(For 2003 data only. Total PDG error now is $\pm 0.13 \times 10^{-7}$)
- **Systematics** will be small.
- **Analysis** far advanced.

Measurement of $K^\pm \rightarrow \pi^\pm \mu^+ \mu^-$



- Event sample of $> 10^3$ $K^\pm \rightarrow \pi^\pm \mu^+ \mu^-$ candidates (SS 1–3).
(Current world sample: 800 events)
- Again: Very small background.



Measurement of $K^\pm \rightarrow \pi^\pm \pi^0 \gamma$

Two amplitudes:

Inner Bremsstrahlung (IB) and Direct Emission (DE).

$$\Gamma \sim \underbrace{\frac{p_\pi^2 \sin^2 \theta}{(m_K/2 - \omega_0)^2}}_{\text{Inner Bremsstrahlung}} \left[1 + \underbrace{2 \frac{m_\pi^2}{m_K^2} W^2 |E| \cos(\theta \pm \delta)}_{\text{Interference}} + \underbrace{\frac{m_\pi^4}{m_K^4} W^4 (|E|^2 + |M|^2)}_{\text{Direct Emission}} \right]$$

Variable $W^2 \propto (p_K q_\gamma)(p_\pi q_\gamma) \propto E_\gamma^{*2} (E_\pi^* - p_\pi^* \cos \theta)$:



Measurement of $K^\pm \rightarrow \pi^\pm \pi^0 \gamma$

■ Previous best measurement: (E787, 2×10^4 events)

Inner Bremsstrahlung (IB): $\text{Br} = (2.75 \pm 0.15) \times 10^{-4}$

Direct Emission (DE): $\text{Br} = (4.7 \pm 0.9) \times 10^{-6}$

Interference term has never been measured.


■ NA48/2 2003 data: (SS 1–3)

More than **200 000 events**.

Backgrounds **< 1%** (tight cuts).

■ Measurements of **branching fraction** and **direct emission component** underway.

⇒ Expect preliminary result soon.



Measurement of $K^\pm \rightarrow \pi^\pm \gamma\gamma$



$K^\pm \rightarrow \pi^\pm \gamma\gamma$: Similar situation as for $K_L \rightarrow \pi^0 \gamma\gamma$:

- $\mathcal{O}(p^2)$ ChPT amplitude vanishes.
- $\mathcal{O}(p^4)$ as function of χ PT parameter \hat{c} : (Ecker, Pich, de Rafael, 1988)

$$\Gamma = (\underbrace{2.80 + 0.87 \hat{c} + 0.14 \hat{c}^2}_{\text{Loop}} + \underbrace{0.23}_{\text{Wess-Zumino}}) \times 10^{-23} \text{ GeV}$$

- $\mathcal{O}(p^6)$ may have contribute by 30–40%. (D'Ambrosio, Portoles, 1996)

Fit of $m_{\gamma\gamma}$ distribution \Rightarrow Information on $\mathcal{O}(p^6)$!





Measurement of $K^\pm \rightarrow \pi^\pm \gamma\gamma$

Experimental problem:

Low- $m_{\gamma\gamma}$ events suppressed at trigger level.

For $m_{\gamma\gamma} > 0.2 \text{ GeV}/c^2$: More than **2000 events** in SS 1–3.

(Compare with 31 events of E787!)



⇒ Promising — but still a lot of work to do!



Rare K_L and K_S Decays

Measurements of $K_L \rightarrow \pi\pi$ Decays

■ Motivation 1: Determine $\Gamma(K_L \rightarrow 3\pi^0)/\Gamma(K_{e3})$

(Large discrepancies in previous measurements \rightarrow Error on $|V_{us}|$)

■ Measure $\Gamma(K_L \rightarrow 2\pi^0)/\Gamma(K_L \rightarrow 3\pi^0)$

■ Measure $\Gamma(K_L \rightarrow \pi^+\pi^-)/\Gamma(K_{e3})$

■ Use $\frac{\Gamma(K_L \rightarrow \pi^+\pi^-)}{\Gamma(K_S \rightarrow \pi^+\pi^-)} / \frac{\Gamma(K_L \rightarrow \pi^0\pi^0)}{\Gamma(K_S \rightarrow \pi^0\pi^0)} = 1 - 6 \times \text{Re}(\epsilon'/\epsilon)$

\Rightarrow NA48, KTeV (2003)

■ Use $\Gamma(K_S \rightarrow \pi^+\pi^-)/\Gamma(K_S \rightarrow \pi^0\pi^0) = 2.236 \pm 0.015$

\Rightarrow KLOE (2002)

\Rightarrow Indirectly measured: $\Gamma(K_L \rightarrow 3\pi^0)/\Gamma(K_{e3})$

■ Motivation 2:

New KTeV result on $K_L \rightarrow \pi^+\pi^-$ disagrees with world average:

KTeV: $|\eta_{+-}| = (2.228 \pm 0.010) \times 10^{-3}$

PDG 2004: $|\eta_{+-}| = (2.288 \pm 0.014) \times 10^{-3}$

\Rightarrow 3.5 σ difference.




Measurements of $K_L \rightarrow \pi\pi$ Decays

$K_L \rightarrow \pi^+\pi^-$:

- More than **60 000 events** in **minimum bias run 99**. (Same as used for K_{e3} .)
- Background tiny ($< 0.2\%$).
- Stat. error $\sigma_{\text{Br}}/\text{Br} \approx 0.4\%$, systematics under study.

$K_L \rightarrow \pi^0\pi^0$:

- Run 2000 with no DCH's. (Same as for $K_L \rightarrow \gamma\gamma$.)
- Expect $> 10^6$ events.



Measurement of $\text{Br}(K_S \rightarrow \pi^+ \pi^- \pi^0)$

- CP-allowed $K_S \rightarrow \pi^+ \pi^- \pi^0$ amplitude suppressed by internal angular momentum of $l = 1$.

- Extraction by using asymmetry in Dalitz plot variable $X = (s_{\pi^-} - s_{\pi^+})/m_{\pi^+}^2 \Rightarrow$ Parameter λ

- Measure $\frac{N^{X>0} - N^{X<0}}{N^{X>0} + N^{X<0}}(t)$


$$\approx D(E) [\text{Re}(2\lambda) \cos(\Delta mt) + \text{Im}(2\lambda) \sin(\Delta mt)] \times e^{-\frac{t}{2}(\frac{1}{\tau_S} - \frac{1}{\tau_S})}$$

- NA48/1 (2002):

\approx 19 million $K^0 \rightarrow \pi^+ \pi^- \pi^0$ events.

- Statistical uncertainties of $\sigma_{\text{Re}(\lambda)} \approx \pm 0.013$, $\sigma_{\text{Im}(\lambda)} \approx \pm 0.010$ competitive to previous experiments.

- Analysis finished, publication soon.



Search for $K_S \rightarrow 3\pi^0$

NA48 result:



$$\text{Re}(\eta_{000}) = -0.002 \pm 0.011_{\text{stat}} \pm 0.015_{\text{sys}}$$

$$\text{Im}(\eta_{000}) = -0.003 \pm 0.013_{\text{stat}} \pm 0.017_{\text{sys}}$$

(CPLEAR: $\text{Re}(\eta_{000}) = 0.18 \pm 0.14 \pm 0.06$
 $\text{Im}(\eta_{000}) = 0.15 \pm 0.20 \pm 0.03$)

If $\text{Re}(\eta_{000}) = \text{Re}(\epsilon)$ (CPT):

$$\text{Im}(\eta_{000}) = 0.000 \pm 0.009_{\text{stat}} \pm 0.013_{\text{sys}}$$



Branching ratio:

$$\text{Br}(K_S \rightarrow 3\pi^0) < 7.4 \times 10^{-7} \quad 90\% \text{ CL}$$

(SND: $\text{Br}(K_S \rightarrow 3\pi^0) < 1.4 \times 10^{-5}$)

$$(\text{Br}(K_S \rightarrow 3\pi^0)|_{\text{CPT}} < 2.3 \times 10^{-7} \quad 90\% \text{ CL})$$

CPT test from $K_S \rightarrow 3\pi^0$

■ Bell-Steinberger relation:

Connects CPT violating phase δ with η parameters via unitarity:

$$(1 + i \tan \phi_{SW}) [\text{Re}(\epsilon) - i \text{Im}(\delta)] = \sum_{\substack{\text{fi nal} \\ \text{states } f}} \alpha_f \quad (\phi_{SW} = \arctan \frac{2 \Delta m}{\Gamma_L - \Gamma_S})$$

■ Largest contributions:

α_f	$10^3 \times \text{Re}(\alpha_f)$	$10^3 \times \text{Im}(\alpha_f)$
$\alpha_{+-} = \eta_{+-} \text{Br}(K_S \rightarrow \pi^+ \pi^-)$	1.146 ± 0.015	1.084 ± 0.016
$\alpha_{00} = \eta_{00} \text{Br}(K_S \rightarrow \pi^0 \pi^0)$	0.511 ± 0.008	0.488 ± 0.008
$\alpha_{+-\gamma} = \eta_{+-\gamma} \text{Br}(K_S \rightarrow \pi^+ \pi^- \gamma)$	0.003 ± 0.000	0.003 ± 0.000
α_{l3}	-0.001 ± 0.007	0.005 ± 0.006
$\alpha_{+-0} = \frac{\tau_S}{\tau_L} \eta_{+-0}^* \text{Br}(K_L \rightarrow \pi^+ \pi^- \pi^0)$	0.000 ± 0.002	0.000 ± 0.002
$\alpha_{000} = \frac{\tau_S}{\tau_L} \eta_{000}^* \text{Br}(K_L \rightarrow 3\pi^0)$	0.029 ± 0.040	-0.026 ± 0.058

- NA48: $\alpha_{000} = (-0.001 \pm 0.007) + i (0.001 \pm 0.008) \times 10^{-3}$
- $\Rightarrow \text{Im}(\delta) = (-0.2 \pm 2.0) \times 10^{-5}$ (was $(2.4 \pm 5.0) \times 10^{-5}$)
- $\Rightarrow m_{K^0} - m_{\overline{K^0}} = (-0.2 \pm 2.8) \times 10^{-19} \text{ GeV}$

■ Rare K^\pm decays:

All measurements with **large statistics** and **low systematics**.

Lots of **on-going work**:

- High precision measurement of K_{e4}^+ .
- Branching fraction and form factors of $K^\pm \rightarrow \pi^\pm e^+ e^-$ and $K^\pm \rightarrow \pi^\pm \mu^+ \mu^-$.
- Advanced analysis on $K^+ \rightarrow \pi^+ \pi^0 \gamma$.
- Work on $K^+ \rightarrow \pi^+ \gamma \gamma$.

■ $K_{L,S}$ decays:

- Precise measurements of $K_L \rightarrow \pi\pi \rightarrow \eta_{+-}, \eta_{00}$ soon.
- Measurements of $K_S \rightarrow \pi^+ \pi^- \pi^0$ and $K_S \rightarrow 3\pi^0$.