

PROSPINO AND LEPTOQUARKS : FROM TEV. TO TEV

Michael Spira (PSI)

I PROSPINO

II Leptoquarks


III Conclusions

in collaboration with W. Beenakker, M. Krämer, T. Plehn and P. Zerwas


I PROSPINO

3 classes of SUSY particle production processes:


- strongly interacting particle pairs:



- weakly interacting particle pairs:




- associated production:




- LO: theoretical uncertainties $\mathcal{O}(100\%) \Rightarrow$ NLO needed

Uncertainties reduced by SUSY-QCD corrections:

— virtual 1 loop contributions



— real contributions due to gluon radiation/crossing



- α_s , PDF: $\overline{\text{MS}}$ scheme [5 active flavors]

- $m_{\tilde{q}}, m_{\tilde{g}}$ on-shell

- double counting: $gq \rightarrow \tilde{g}\tilde{q} \rightarrow \tilde{g}\tilde{\chi}q$ [if $m_{\tilde{q}} > m_{\tilde{\chi}}$]

$$\frac{d\hat{\sigma}_{res}}{dM^2} = \hat{\sigma}(gq \rightarrow \tilde{g}\tilde{q}) BR(\tilde{q} \rightarrow \tilde{\chi}q) \underbrace{\frac{m_{\tilde{q}}\Gamma_{\tilde{q}}/\pi}{(M^2 - m_{\tilde{q}}^2)^2 + m_{\tilde{q}}^2\Gamma_{\tilde{q}}^2}}_{\rightarrow \delta(M^2 - m_{\tilde{q}}^2)}$$

to be subtracted

[analogous subtraction for $\tilde{q} \rightarrow \tilde{g}q$]

central scale: $K = \sigma_{NLO}/\sigma_{LO} \sim 1.1 - 1.2$

F

$\frac{1}{2}m < \mu_R = \mu_F < 2m: \delta\sigma \lesssim \pm 5 - 10\%$

F

\Rightarrow NLO corrections lead to reliable predictions



[now agreement with Berger, Klasen, Tait]

- $pp \rightarrow \tilde{q}\tilde{\chi}$: first very preliminary results



F

Beenakker, Krämer, Plehn, S., Zerwas

[resonance subtractions: $\tilde{q} \rightarrow \tilde{\chi}q, \tilde{g} \rightarrow \tilde{q}\bar{q}$]




Beenakker, Krämer,
 Plehn, S., Zerwas



Beenakker, Krämer,
 Plehn, S., Zerwas

VERY PRELIMINARY
LHC



Beenakker, Krämer, Plehn, S., Zerwas

PROSPINO

- $\tilde{g}\tilde{g}, \tilde{q}\tilde{q}, \tilde{q}\tilde{\bar{q}}, \tilde{g}\tilde{g}, \tilde{t}\tilde{\bar{t}}$ production:



$$\sigma_{tot}, \quad \frac{d^2\sigma}{dp_T dy} \text{ @ } NLO$$

Beenakker,...

[\tilde{b} with large mass splitting: $\rightarrow \tilde{t}\tilde{\bar{t}}$ program]

- $\tilde{\chi}\tilde{\chi}, \tilde{\ell}\tilde{\bar{\ell}}, \tilde{g}\tilde{\chi}$ production: σ_{tot} @ NLO added
- $\tilde{q}\tilde{\chi}$ production: first NLO results soon...
- distributions: coming soon...
- new version 2.0: extension to $\tilde{\chi}\tilde{\chi}, \tilde{g}\tilde{\chi}, \tilde{q}\tilde{\chi}$ [LO]

PROSPINO



Beenakker, Krämer, Plehn, S., Zerwas

II LEPTOQUARKS


- leptoquarks predicted by many exotic extension of the SM [compositeness] Buchmüller, Rückl, Wyler

- leptoquarks carry lepton and baryon number



- two basic types: scalar [spin 0], vector [spin 1] (non-renormalizeable)

- novel Yukawa couplings small for first two generations
⇒ negligible for hadroproduction

- relevant for main decay modes: $LQ \rightarrow \ell q, \nu_{\ell} q$



- production: $gg, q\bar{q} \rightarrow LQ \bar{LQ}$




- Leading order: $[\beta = \sqrt{1 - 4M_{LQ}^2/\hat{s}}]$

$$\hat{\sigma}_{gg} = \frac{\alpha_s^2 \pi}{96 \hat{s}} \left[\beta(41 - 31\beta^2) + (18\beta^2 - \beta^4 - 17) \log \frac{1 + \beta}{1 - \beta} \right]$$

$$\hat{\sigma}_{q\bar{q}} = \frac{\alpha_s^2 \pi}{\hat{s}} \frac{2}{27} \beta^3$$


- production model-independent [$M_{LQ} < \Lambda$]
- Tevatron: $q\bar{q}$ channel dominant (75–85%), gg sizeable (15–25%)
- LHC: gg channel dominant (70–95%), $q\bar{q}$ sizeable (5–30%)

Comparison with Other Colliders



λ dependence

1st: $M_{LQ} \gtrsim 250$ GeV
 2nd: $M_{LQ} \gtrsim 240$ GeV
 reach @ Run II: ~ 300 GeV




Simona Rolli, TeV4LHC



QCD CORRECTIONS


- LO: large scale dependence: \implies NLO needed

(i) Virtual Corrections



- ultraviolet, infrared and collinear divergences: dimensional regularization in $n = 4 - 2\epsilon$ dimensions
- α_s : $\overline{\text{MS}}$ scheme [5 flavours], M_{LQ} : on-shell



(ii) Real Corrections





- infrared and collinear singularities cancel against virtual corrections and counter terms of PDFs [mass factorization]
- PDF: $\overline{\text{MS}}$ scheme [5 flavours]

(iii) Cross Checks

- Scalar leptoquark pair production extracted from $\tilde{q}\tilde{q}$ and $\tilde{t}\tilde{t}$ pair production for $m_{\tilde{g}} \rightarrow \infty$ [contributions from quartic scalar couplings subtracted]
- \Rightarrow full agreement









Krämer, Plehn, S., Zerwas

III CONCLUSIONS

- SUSY particle searches at Tevatron/LHC belong to major endeavours
- sensitive up to $m_{\tilde{q},\tilde{g}} \sim 400$ GeV (Tev.), $\sim 2 - 3$ TeV (LHC)
- most (SUSY-)QCD corrections known \Rightarrow large corrections
remaining theoretical uncertainties: $\sim 100\% \longrightarrow \lesssim 10 - 15\%$
- significantly increased mass reaches [Tev.: $\lesssim 30$ GeV, LHC: $\lesssim 50$ GeV]
- program package available with these corrections: **PROSPINO 2.0**
<http://pheno.physics.wisc.edu/~plehn/prospino/prospino.html>
<http://people.web.psi.ch/spira/>
- Scalar leptoquark pair production: QCD corrections $\sim (20 - 90)\%$
 \Rightarrow increased mass reach [Tev.: ~ 15 GeV, LHC: ~ 100 GeV]
- sensitive up to ~ 300 GeV (Tev.), ~ 1.6 TeV (LHC)
- strong reduction of scale dependence: $\Delta \lesssim 10 - 15\%$
- FORTRAN code added to PROSPINO 2.0


BACKUP SLIDES







$$m_{\tilde{q}} = 250 \text{ GeV}$$

$$m_{\tilde{g}} = 300 \text{ GeV}$$

Beenakker, Höpker, S., Zerwas







Beenakker, Höpker, S., Zerwas



$$m_{\tilde{q}} = 250 \text{ GeV}$$

$$m_{\tilde{g}} = 300 \text{ GeV}$$


Beenakker, Höpker, S., Zerwas



$$m_{\tilde{q}} = 600 \text{ GeV}$$




$$m_{\tilde{g}} = 500 \text{ GeV}$$

Beenakker, Höpker, S., Zerwas




$$m_{\tilde{q}} = 600 \text{ GeV}$$



$$m_{\tilde{g}} = 500 \text{ GeV}$$







Beenakker, Höpker, S., Zerwas



Beenakker, Höpker, S., Zerwas





Beenakker, Krämer, Plehn, S., Zerwas



$$m_{\tilde{q}} = 250 \text{ GeV}$$

$$m_{\tilde{g}} = 300 \text{ GeV}$$



Beenakker, Klasen, Krämer, Plehn, S., Zerwas